

A matrix second-order negative Ablowitz–Kaup–Newell–Segur flow and its Darboux transformation

Wen-Xiu Ma 

*Department of Mathematics, Zhejiang Normal University
Jinhua 321004, P. R. China*

*Research Center of Astrophysics and Cosmology
Khazar University, 41 Mehseti Street
Baku 1096, Azerbaijan*

*Department of Mathematics and Statistics, University of South Florida
Tampa, FL 33620-5700, USA*

*Material Science Innovation and Modelling
North-West University, Mafikeng Campus, Private Bag X2046
Mmabatho 2735, South Africa
maux@cas.usf.edu*

Received 29 July 2025

Accepted 22 October 2025

Published 22 November 2025

Negative-order integrable flows have significant applications in areas such as fluid dynamics, nonlinear optics, and quantum field theory, particularly in the study of soliton behavior and wave interactions. In this paper, we study a matrix second-order negative flow within the Ablowitz–Kaup–Newell–Segur (AKNS) hierarchy, formulated through a Lax pair whose temporal part involves a second-order pole in the spectral parameter. This framework facilitates the analysis of shallow water wave dynamics, including the emergence of lump and soliton solutions. The corresponding Darboux transformation is constructed within the general AKNS-type framework. Starting from a seed solution, a class of explicit solutions is generated through a single application of the Darboux transformation.

Keywords: Lax pair; matrix AKNS flows; Darboux transformation.

Mathematics Subject Classification 2020: 35Q51, 35Q58, 37K40

PACS: 02.30.Ik, 05.45.Yv

1. Introduction

Negative-order flows constitute a fascinating and increasingly significant area within the theory of integrable models. These flows are deeply connected to rich mathematical structures, such as infinite-dimensional symmetries, recursion operators, and algebraic curves, as well as to a variety of physical applications, particularly in fluid dynamics and nonlinear optics (see, e.g. [1–3]). A prominent example is the

Camassa–Holm equation [4], an integrable model with a bi-Hamiltonian structure that describes unidirectional shallow water waves and supports peakon solutions. The study of negative-order flows enhances our understanding of nonlinear wave phenomena, inverse dynamics, and soliton interactions, extending beyond the scope of classical integrable equations.

The Darboux transformation is a powerful method used to generate new solutions from known ones for integrable partial differential equations, especially in soliton theory. It is typically applied to linear spectral problems, known as a Lax pair, associated with nonlinear integrable equations.

Originally introduced by Gaston Darboux in the 19th century in the context of second-order linear differential equations, the transformation was later adapted and extensively developed in the 20th century for integrable systems. In particular, its application to the Korteweg–de Vries hierarchy and other soliton equations has played a pivotal role in modern soliton theory. A comprehensive treatment of the Darboux transformation in integrable systems is provided in the monograph by Matveev and Salle [5]. Further developments, including generalizations to matrix, noncommutative, super-symmetric, and higher-dimensional settings, can be found in [6–8].

At its core, the Darboux transformation constructs a new potential (or solution) by applying an algebraic transformation to the Lax pair of an integrable system. This process preserves the integrability of the original system and allows the recursive generation of multi-soliton, rational, and rogue wave solutions (see, e.g. [9, 10]).

Let us now recall the zero-curvature representation and the formulation of Darboux transformations within this framework. A system of partial differential equations

$$u_t = K(u) = K(x, t, u, u_x, \dots) \quad (1.1)$$

is said to admit a zero-curvature equation representation, if it arises from the compatibility condition of a pair of linear equations:

$$U_t - V_x + [U, V] = 0, \quad (1.2)$$

where U and V are square matrices, known as a Lax pair, depending on the solution u , its derivatives, and the spectral parameter λ . These matrices typically lie in a matrix loop algebra [11]. The zero-curvature equation is equivalent to the compatibility of the linear system

$$\phi_x = U\phi = U(u, \lambda)\phi, \quad \phi_t = V\phi = V(u, \lambda)\phi, \quad (1.3)$$

where ϕ is the eigenfunction vector and λ is the spectral parameter.

A transformation of $\phi' = D\phi$ and $u' = u'(u)$, where $D = D(u, \lambda)$ is a square matrix, is called a Darboux transformation of the spectral problems (1.3), if ϕ' satisfies the same type spectral problems:

$$\phi'_x = U'\phi' = U(u', \lambda)\phi', \quad \phi'_t = V'\phi' = V(u', \lambda)\phi'. \quad (1.4)$$

The matrix D is referred to as the Darboux matrix. The transformed Lax pair (U', V') must satisfy the gauge-type conditions:

$$U'D = DU + D_x, \quad V'D = DV + D_t. \quad (1.5)$$

When U and V are $N \times N$ matrices, a common ansatz for a first-order Darboux matrix is

$$D(\lambda) = \lambda I_N - S, \quad (1.6)$$

where I_N is the identity matrix of order N and S is an $N \times N$ matrix independent of λ . To construct S , one chooses N eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_N$ and their corresponding eigenfunctions solving

$$\phi_x^{[j]} = U(u, \lambda_j)\phi^{[j]}, \quad \phi_t^{[j]} = V(u, \lambda_j)\phi^{[j]}, \quad 1 \leq j \leq N, \quad (1.7)$$

with u being a fixed solution to (1.1). Then, the matrix S is given by [6, 12]:

$$S = HAH^{-1}, \quad H = (\phi^{[1]}, \dots, \phi^{[N]}), \quad A = \text{diag}(\lambda_1, \dots, \lambda_N). \quad (1.8)$$

We summarize the Darboux transformation as follows:

Darboux transformation:		
$D = \lambda I_N - H\Lambda H^{-1}$		
Lax pair: (U, V)	$\implies$	New Lax pair: (U', V')
Solution: $u_t = K(u)$	$\implies$	New solution: $u'_t = K(u')$

In this work, we begin with the matrix Ablowitz–Kaup–Newell–Segur (AKNS) spectral problem and derive a matrix-valued second-order negative AKNS flow by introducing a Lax pair involving a second-order pole in the spectral parameter. This flow is a natural extension of the Camassa–Holm type equations associated with the AKNS spectral problem. Based on this Lax representation, we construct the corresponding Darboux transformation within the AKNS framework. By applying this transformation to a seed solution, we generate a class of explicit solutions for the proposed matrix second-order negative AKNS system.

2. Lax Pair and An Associated Integrable System

Let m, n be two natural numbers. We consider the AKNS matrix spectral matrix of the form [13]:

$$U = i\lambda\Lambda + Q, \quad \Lambda = \begin{bmatrix} -I_m & 0 \\ 0 & I_n \end{bmatrix}, \quad Q = \begin{bmatrix} 0 & q \\ r & 0 \end{bmatrix}, \quad (2.1)$$

where I_k denotes the identity matrix of order k , and q and r are the two potential matrices given by

$$q = (q_{jk})_{m \times n}, \quad r = (r_{kj})_{n \times m}. \quad (2.2)$$

We formulate the Lax matrix as follows:

$$V = \frac{1}{1 - 4\lambda^2} W, \quad W = \lambda W_1 + W_2 \quad (2.3)$$

with

$$W_1 = i \begin{bmatrix} u_1 & 0 \\ 0 & u_2 \end{bmatrix} \Lambda - 2i\Lambda Q_t, \quad W_2 = -\frac{1}{2} \begin{bmatrix} v_1 & f \\ g & v_2 \end{bmatrix} \Lambda - Q_{tx}, \quad (2.4)$$

where u_1, v_1 and u_2, v_2 are square potential matrices of orders m and n , respectively, and the off-diagonal blocks f and g are defined as

$$f = -u_1 q - q u_2, \quad g = r u_1 + u_2 r. \quad (2.5)$$

Note that

$$\begin{bmatrix} 0 & f \\ g & 0 \end{bmatrix} = [\Delta\Lambda, Q], \quad \Delta = \begin{bmatrix} u_1 & 0 \\ 0 & u_2 \end{bmatrix}.$$

We present the Lax matrix V in this form to simplify the resulting model equations.

The zero-curvature equation

$$U_t - V_x + [U, V] = 0 \quad (2.6)$$

is equivalent to

$$(1 - 4\lambda^2)U_t - W_x + [U, W] = 0.$$

By matching powers of λ , we obtain the system

$$\begin{cases} -4Q_t + i[\Lambda, W_1] = 0, \\ -W_{1,x} + i[\Lambda, W_2] + [Q, W_1] = 0, \\ Q_t - W_{2,x} + [Q, W_2] = 0. \end{cases} \quad (2.7)$$

The first equation is automatically satisfied. From the second and third equations, we derive the following system:

$$\begin{cases} q_t + q_{txx} - \frac{1}{2}(u_1 q + q u_2)_x - \frac{1}{2}(v_1 q + q v_2) = 0, \\ r_t + r_{txx} - \frac{1}{2}(r u_1 + u_2 r)_x + \frac{1}{2}(r v_1 + v_2 r) = 0, \\ v_{1,x} + 2q r_{tx} - 2q_{tx} r = 0, \\ v_{2,x} - 2r q_{tx} + 2r_{tx} q = 0, \\ u_{1,x} - 2(qr)_t = 0, \\ u_{2,x} - 2(rq)_t = 0. \end{cases} \quad (2.8)$$

The first four equations come from the third equation in (2.7), and the last two come from the second equation. Because the spectral parameter λ appears with a second-order term in the denominator of V , this system corresponds to a second-order negative AKNS flow. It is actually a special case of the system presented in [12], with

$$\begin{cases} n = 2, & p_1 = x, & x_2 = t, & A_1 = -i\lambda\Lambda - Q, \\ a_2 = 1 - 4\lambda^2, & A_2 = \lambda W_1 + W_2. \end{cases} \quad (2.9)$$

Note that all functions involved in U and V above, such as q and r , are functions of x and t only, and therefore do not depend on x_1 and p_2 .

When we take the following in our construction:

$$m = n = 1, \quad u_1 = u_2 = w, \quad v_1 = v_2 = v, \quad (2.10)$$

the Lax operator V reduces to the scalar form given in [14], and the system (2.8) simplifies to

$$\begin{cases} q_t + q_{txx} - (wq)_x - vq = 0, \\ r_t + r_{txx} - (wr)_x + vr = 0, \\ v_x - 2(rq_{tx} - r_{tx}q) = 0, \\ w_x - 2(qr)_t = 0. \end{cases} \quad (2.11)$$

This is exactly the Nurshuak–Tolkynay–Myrzakulov-II equation as introduced in [14]. There are other negative integrable flows associated with the AKNS spectral problem (see, e.g. [15–19]).

3. Darboux Transformation and Explicit Solutions

3.1. Compatibility conditions

We assume the Darboux matrix takes the form

$$D(\lambda) = \lambda I_{m+n} - S. \quad (3.1)$$

W. X. Ma

Then, D must satisfy the spatial compatibility condition:

$$U'D = DU + D_x, \quad (3.2)$$

where U is given by (2.1) and

$$U' = i\lambda\Lambda + Q'. \quad (3.3)$$

This leads to

$$(i\lambda\Lambda + Q')(\lambda I_{m+n} - S) = (\lambda I_{m+n} - S)(i\lambda\Lambda + Q) - S_x$$

from which we deduce

$$Q' = Q + i[\Lambda, S] \quad (3.4)$$

and

$$S_x = Q'S - SQ = [Q + i\Lambda S, S]. \quad (3.5)$$

Next, we consider the temporal compatibility condition:

$$V'D = DV + D_t, \quad (3.6)$$

where V is defined by (2.3) and

$$V' = \frac{1}{1-4\lambda^2}W', \quad W' = \lambda W'_1 + W'_2. \quad (3.7)$$

This implies

$$W'(\lambda I_{m+n} - S) = (\lambda I_{m+n} - S)W - (1-4\lambda^2)S_t.$$

Comparing coefficients of powers of λ , we obtain

$$\begin{cases} W'_1 = W_1 + 4S_t, \\ W'_2 = W_2 + W'_1 S - SW_1, \end{cases} \quad (3.8)$$

and the evolution of S satisfies:

$$S_t = W'_2 S - SW_2. \quad (3.9)$$

3.2. Formulation of the Darboux matrix

Following the general framework (cf. [12]), we define the matrix S as

$$S = HAH^{-1}, \quad H = (\phi^{[1]}, \dots, \phi^{[m+n]}), \quad A = \text{diag}(\lambda_1, \dots, \lambda_{m+n}), \quad (3.10)$$

where each column vector $\phi^{[j]}$ satisfies the eigenvalue problems:

$$\phi_x^{[j]} = U(u, \lambda_j) \phi^{[j]}, \quad \phi_t^{[j]} = V(u, \lambda_j) \phi^{[j]}, \quad 1 \leq j \leq m+n. \quad (3.11)$$

From this, we compute

$$H_x = i\Lambda HA + QH, \quad H_t = W_1 HB_1 + W_2 HB_2, \quad (3.12)$$

where the diagonal matrices B_1 and B_2 are given by

$$\begin{aligned} B_1 &= \text{diag} \left(\frac{\lambda_1}{1-4\lambda_1^2}, \dots, \frac{\lambda_{m+n}}{1-4\lambda_{m+n}^2} \right), \\ B_2 &= \text{diag} \left(\frac{1}{1-4\lambda_1^2}, \dots, \frac{1}{1-4\lambda_{m+n}^2} \right). \end{aligned} \quad (3.13)$$

Using (3.12), we compute the derivatives S_x and S_t , and verify that both the spatial and temporal compatibility conditions (3.5) and (3.9) are satisfied.

The spatial derivative yields:

$$\begin{aligned} S_x &= H_x AH^{-1} - HA(H^{-1}H_x H^{-1}) \\ &= i\Lambda HA^2 H^{-1} + QHAH^{-1} - HAH^{-1}(i\Lambda HA + QH)H^{-1} \\ &= i\Lambda HA^2 H^{-1} + QHAH^{-1} - iHAH^{-1}\Lambda HAH^{-1} - HAH^{-1}Q \\ &= QS - SQ + i\Lambda S^2 - i\Lambda S, \end{aligned}$$

which confirms that the spatial compatibility condition (3.5) holds.

To verify the temporal condition, we compute

$$\begin{aligned} S_t &= H_t AH^{-1} - HA(H^{-1}H_t H^{-1}) \\ &= W_1 HB_1 AH^{-1} + W_2 HB_2 AH^{-1} - HAH^{-1}W_1 HB_1 H^{-1} \\ &\quad - HAH^{-1}W_2 HB_2 H^{-1}. \end{aligned}$$

Based on this, we then define

$$\begin{aligned} h_1(W_1) + h_2(W_2) &:= W_2' S - SW_2 - S_t = [W_2 + (W_1 + 4S_t)S - SW_1]S \\ &\quad - SW_2 - S_t, \end{aligned} \quad (3.14)$$

where h_1 collects all terms containing W_1 , and h_2 those containing W_2 . Using the identities

$$4B_1 A^2 = B_1 - A, \quad 4B_2 A^2 = B_2 - I_{m+n}, \quad (3.15)$$

we can compute that

$$\begin{aligned} h_1(W_1) &\equiv W_1 HA^2 H^{-1} + 4W_1 HB_1 A^3 H^{-1} - 4HAH^{-1}W_1 HB_1 A^2 H^{-1} \\ &\quad - HAH^{-1}W_1 HAH^{-1} - W_1 HB_1 AH^{-1} + HAH^{-1}W_1 HB_1 H^{-1} \\ &= W_1 H(A + 4B_1 A^2 - B_1)AH^{-1} - HAH^{-1}W_1 H(4B_1 A^2 + A - B_1)H^{-1} \\ &= 0 \end{aligned}$$

W. X. Ma

and

$$\begin{aligned}
h_2(W_2) &\equiv W_2 H A H^{-1} + 4W_2 H B_2 A^3 H^{-1} - 4H A H^{-1} W_2 H B_2 A^2 H^{-1} \\
&\quad - H A H^{-1} W_2 - W_2 H B_2 A H^{-1} + H A H^{-1} W_2 H B_2 H^{-1} \\
&= W_2 H (I_{m+n} + 4B_2 A^2 - B_2) A H^{-1} \\
&\quad - H A H^{-1} W_2 H (4B_2 A^2 + I_{m+n} - B_2) H^{-1} \\
&= 0.
\end{aligned}$$

These computations confirm the validity of the temporal compatibility condition (3.9).

3.3. The Darboux transformation

Therefore, the Darboux transformation is given by

$$\phi' = (\lambda I_{m+n} - S)\phi, \quad U' = i\lambda\Lambda + Q', \quad V' = \frac{1}{1 - 4\lambda^2}(\lambda W'_1 + W'_2) \quad (3.16)$$

with

$$Q' = Q + i[\Lambda, S], \quad W'_1 = W_1 + 4S_t, \quad W'_2 = W_2 + W'_1 S - S W_1, \quad (3.17)$$

where the matrix S is defined by (3.10). Observing the structures of U , W_1 , and W_2 in (2.1) and (2.4), this transformation yields the following expressions for the transformed quantities:

$$\begin{cases} q' = q + i[\Lambda, S]_{12}, & r' = r + i[\Lambda, S]_{21}, \\ u'_1 = u_1 + 4iS_{11,t}, & u'_2 = u_2 - 4iS_{22,t}, \\ v'_1 = v_1 + 2[W_1, S]_{11} + 8(S_t S)_{11}, & v'_2 = v_2 - 2[W_1, S]_{22} - 8(S_t S)_{22}. \end{cases} \quad (3.18)$$

Here, a matrix A of order $m + n$ is partitioned into block form as

$$M = \left[\begin{array}{c|c} M_{11} & M_{12} \\ \hline M_{21} & M_{22} \end{array} \right] = \left[\begin{array}{c|c} (M_{11})_{m \times m} & (M_{12})_{m \times n} \\ \hline (M_{21})_{n \times m} & (M_{22})_{n \times n} \end{array} \right].$$

3.4. Application: Explicit solutions

Consider a seed solution

$$q = r = 0, \quad u_1 = u_{1,0}, \quad u_2 = u_{2,0}, \quad v_1 = v_{1,0}, \quad v_2 = v_{2,0}, \quad (3.19)$$

where $u_{1,0}$, $u_{2,0}$, $v_{1,0}$, and $v_{2,0}$, are constant matrices. The fundamental eigenfunctions

$$\phi^{[j]} = (\phi_1^{[j]T}, \phi_2^{[j]T})^T \quad (3.20)$$

take the form

$$\begin{cases} \phi_1^{[j]} = \exp \left(i\lambda_j I_m x - \frac{1}{1-4\lambda_j^2} \left(i\lambda_j u_{1,0} - \frac{1}{2}v_{1,0} \right) t \right) \mu_1^{[j]}, \\ \phi_2^{[j]} = \exp \left(-i\lambda_j I_n x + \frac{1}{1-4\lambda_j^2} \left(i\lambda_j u_{2,0} - \frac{1}{2}v_{2,0} \right) t \right) \mu_2^{[j]}, \end{cases} \quad 1 \leq j \leq m+n, \quad (3.21)$$

where $\mu_1^{[j]} \in \mathbb{C}^m$ and $\mu_2^{[j]} \in \mathbb{C}^n$ are arbitrary constant column vectors. Due to $W_1 = \text{diag}(-iu_{1,0}, iu_{2,0})$, we have $[W_1, S]_{11} = 0$ and $[W_1, S]_{22} = 0$, and the resulting explicit solution is then given by

$$\begin{cases} q' = i[\Lambda, S]_{12}, & r' = i[\Lambda, S]_{21}, \\ u'_1 = u_{1,0} + 4iS_{11,t}, & u'_2 = u_{2,0} - 4iS_{22,t}, \\ v'_1 = v_{1,0} + 8(S_t S)_{11}, & v'_2 = v_{2,0} - 8(S_t S)_{22}. \end{cases} \quad (3.22)$$

where $S = HAH^{-1}$ with $H = (\phi^{[1]}, \dots, \phi^{[m+n]})$ and $A = \text{diag}(\lambda_1, \dots, \lambda_{m+n})$.

4. Conclusions

Starting from a specific matrix Lax pair, we have constructed a matrix second-order negative AKNS flow and its associated Darboux transformation, along with a class of explicit solutions. In the scalar case, this construction recovers the Nurshuak–Tolkynay–Myrzakulov-II equation introduced in [14]. Additional developments on Darboux transformations for negative integrable flows can be found in the literature (see, e.g. [20–23]).

Constrained Lax pairs are of particular interest, especially when reductions are imposed on the general potential matrix U (see, e.g. [24, 25]). A natural question then arises: how can one construct Darboux transformations for such constrained matrix systems, given their reduced Lax pairs? This question becomes even more intriguing in the broader context where the Lax operator V contains higher-order poles in its dependence on the spectral parameter λ . A compelling open problem is thus to determine the structure of Darboux transformations in this more general negative-order setting, where both constraints and nontrivial spectral singularities come into play.

Acknowledgments

The work was supported in part by the Ministry of Science and Technology of China (Nos. G2021016032L and G2023016011L) and the National Natural Science Foundation of China (Nos. 12271488 and 11975145).

ORCID

Wen-Xiu Ma  <https://orcid.org/0000-0001-5309-1493>

References

- [1] R. S. Johnson, Camassa–Holm, Korteweg–de Vries and related models for water waves, *J. Fluid Mech.* **455** (2002) 63–82.
- [2] H. Aratyn, J. F. Gomes and A. H. Zimerman, On negative flows of the AKNS hierarchy and a class of deformations of a bihamiltonian structure of hydrodynamic type, *J. Phys. A: Math. Gen.* **39** (2006) 1099–1114.
- [3] Z. J. Qiao and E. G. Fan, Negative-order Korteweg–de Vries equations, *Phys. Rev. E* **86** (2012) 016601.
- [4] R. Camassa and D. D. Holm, An integrable shallow water equation with peaked solitons, *Phys. Rev. Lett.* **71** (1993) 1661–1664.
- [5] V. B. Matveev and M. A. Salle, *Darboux Transformation and Solitons* (Springer, Berlin, 1991).
- [6] C. H. Gu, H. S. Hu and Z. X. Zhou, *Darboux Transformations in Integrable Systems* (Springer, Dordrecht, 2005).
- [7] E. V. Doktorov and S. B. Leble, *A Dressing Method in Mathematical Physics* (Springer, Dordrecht, 2007).
- [8] W. X. Ma and Y. J. Zhang, Darboux transformations of integrable couplings and applications, *Rev. Math. Phys.* **30** (2018) 1850003.
- [9] L. Cheng, Y. Zhang and W. X. Ma, An extended (2+1)-dimensional modified Korteweg–de Vries–Calogero–Bogoyavlenskii–Schiff equation: Lax pair and Darboux transformation, *Comm. Theor. Phys.* **77** (2025) 035002.
- [10] X. Lü, L. L. Zhang and W. X. Ma, Oceanic shallow-water description with (2+1)-dimensional generalized variable-coefficient Hirota–Satsuma–Ito equation: Painlevé analysis, soliton solutions, and lump solutions, *Phys. Fluids* **36** (2024) 064110.
- [11] W. X. Ma and M. Chen, Hamiltonian and quasi-Hamiltonian structures associated with semi-direct sums of Lie algebras, *J. Phys. A: Math. Gen.* **39** (2006) 10787–10801.
- [12] W. X. Ma, Darboux transformations for a Lax integrable system in $2n$ -dimensions, *Lett. Math. Phys.* **39** (1997) 33–49.
- [13] M. J. Ablowitz, D. J. Kaup, A. C. Newell and H. Segur, The inverse scattering transform-Fourier analysis for nonlinear problems, *Stud. Appl. Math.* **53** (1974) 249–315.
- [14] Z. Sagidullayeva, G. Nugmanova, R. Myrzakulov and N. Serikbayev, Integrable Kuralay equations: Geometry, solutions and generalizations, arXiv:2206.05348v2.
- [15] D. J. Zhang, J. Ji and S. L. Zhao, Soliton scattering with amplitude changes of a negative order AKNS equation, *Phys. D* **238** (2009) 2361–2367.
- [16] J. Ji, J. B. Zhang and D. J. Zhang, Soliton solutions for a negative order AKNS equation hierarchy, *Commun. Theor. Phys.* **52** (2009) 395–397.
- [17] Y. Q. Yao and Y. B. Zeng, Coupled short pulse hierarchy and its Hamiltonian structure, *J. Phys. Soc. Jpn.* **80** (2011) 064004.
- [18] M. Gürses and A. Pekcan, (2+1)-dimensional AKNS(- N) systems II, *Commun. Non-linear Sci. Numer. Simul.* **97** (2021) 105736.
- [19] X. G. Geng, J. Wang and B. Xue, Riemann-Hilbert approach and N -soliton solutions for a negative matrix AKNS system with a Hermitian symmetric space, *Wave Motion* **108** (2022) 102838.
- [20] C. Chen and Z. X. Zhou, Darboux transformation and exact solutions of the Myrzakulov-I equation, *Chin. Phys. Lett.* **26** (2009) 080504.
- [21] H. Wajahat and A. Riaz, Darboux transformation for a negative order AKNS equation, *Commun. Theor. Phys.* **71** (2019) 912–920.

- [22] S. L. Huang and H. M. Li, Darboux transformations and soliton-like solutions of a new system associated with the negative AKNS system, *Adv. Math. Phys.* **2022** (2022) 3276320.
- [23] F. Müller-Hoissen, A vectorial binary Darboux transformation for the first member of the negative part of the AKNS hierarchy, *J. Phys. A: Math. Theor.* **56** (2023) 125701.
- [24] W. X. Ma, Integrable matrix nonlinear Schrödinger equations with reduced Lax pairs of AKNS type, *Appl. Math. Lett.* **168** (2025) 109574.
- [25] W. X. Ma, Integrable matrix modified Korteweg-de Vries equations derived from reducing AKNS Lax pairs, *Rom. J. Phys.* **70** (2025) 110.