

Wen-Xiu Ma*

Lump-Type Solutions to the (3+1)-Dimensional Jimbo-Miwa Equation

DOI 10.1515/ijnsns-2015-0050

Received April 20, 2015; accepted October 7, 2016

Abstract: Taking advantage of the Hirota bilinear form, four classes of lump-type solutions to the (3+1)-dimensional Jimbo-Miwa equation are presented through symbolic computation with Maple. Special choices of the involved parameters guaranteeing analyticity of the fourth solution are given, together with two particular lump-type solutions.

Keywords: Hirota bilinear form, Jimbo-Miwa equation, Lump-type solution

PACS® (2010). 02.30.Ik

MSC® (2010). 35Q51, 35Q53, 37K40

1 Introduction

Nonlinear integrable equations can be transformed into Hirota bilinear equations, and such examples of equations include the Korteweg-de Vries (KdV) equation, the Boussinesq equation, the Kadomtsev–Petviashvili (KP) equation and the Toda lattice equation [1]. All these integrable equations possess exponentially localized solutions – soliton solutions [2]. Hirota bilinear forms play a crucial role in generating soliton solutions, though some intelligent guesswork is often needed [3].

Recently, there has been a renewed and growing interest in rational solutions to nonlinear partial differential equations (see, e.g., [4, 5]). Particularly important are rationally localized solutions, called lump solutions, and examples of lump solutions are found for many integrable equations such as the KP equation I [6], the three-dimensional three-wave resonant interaction [7], the B-KP equation [8], the Davey–Stewartson equation II [9] and the Ishimori-I equation [10]. The KP equation I of the form

$$(u_t + 6uu_x + u_{xxx})_x - 3u_{yy} = 0 \quad (1)$$

possesses the following lump solution [6]:

$$u = 4 \frac{-[x + ay + 3(a^2 - b^2)t]^2 + b^2(y + 6at)^2 + 1/b^2}{\{[x + ay + 3(a^2 - b^2)t]^2 + b^2(y + 6at)^2 + 1/b^2\}^2}, \quad (2)$$

where a and b are real free parameters. Rogue wave solutions, which draw a big attention of mathematicians and physicists worldwide, are a particularly interesting kind of lump or lump-type solutions, and such solutions, usually with rational function amplitudes, could be used to describe significant nonlinear wave phenomena in both oceanography [11] and nonlinear optics [12]. It is natural and interesting to search for lump or lump-type solutions to nonlinear partial differential equations, on the basis of Hirota bilinear forms.

General rational solutions to nonlinear integrable equations have been considered within the Wronskian formulation, the Casoratian formulation and the Pfaffian formulation (see [1, 2]). The KdV equation and the Boussinesq equation in (1+1)-dimensions, the KP equation in (2+1)-dimensions and the Toda lattice equation in (0+1)-dimensions are such typical examples (see, e.g., [13–16]). Several attempts have also been made to search for rational solutions to the non-integrable (3+1)-dimensional KP I [17, 18] and KP II [19] by direct approaches such as the tanh-function method and the $\frac{G'}{G}$ -expansion method (see, e.g., [20, 21]). Rational solutions to the (3+1)-dimensional KP II are linked to the good Boussinesq equation by a transformation of dependent variables [19]. Moreover, bilinear Bäcklund transformations are used to construct rational solutions to (3+1)-dimensional generalized KP equations (see, e.g., [22]), and there is some direct search for rational solutions to generalized bilinear equations (see, e.g., [23]), formulated in terms of generalized bilinear derivatives [24].

In this paper, we would like to focus on the (3+1)-dimensional Jimbo-Miwa equation and present four classes of its lump-type solutions by symbolic computation with Maple. The (3+1)-dimensional Jimbo-Miwa equation has a Hirota bilinear form, and so, we will do a search for positive quadratic function solutions to the corresponding (3+1)-dimensional bilinear Jimbo-Miwa equation. The obtained quadratic function solutions contain a set

*Corresponding author: Wen-Xiu Ma, Department of Mathematics, Zhejiang Normal University, Jinhua 321004, PR China; College of Mathematics and Physics, Shanghai University of Electric Power, Shanghai 200090, PR China; Department of Mathematics and Statistics, University of South Florida, Tampa, FL 33620, USA, E-mail: mawx@cas.usf.edu

of free parameters, and taking special choices of parameters involved, particular lump-type solutions will be generated from the fourth quadratic function solution. A few concluding remarks are given finally at the end of the paper.

2 Lump-type solutions to the Jimbo-Miwa equation

2.1 The Jimbo-Miwa equation

The (3+1)-dimensional Jimbo-Miwa equation reads [25]

$$P_{JM}(u) := u_{xxxy} + 3u_y u_{xx} + 3u_x u_{xy} + 2u_{yt} - 3u_{xz} = 0, \quad (3)$$

called the Jimbo-Miwa equation in [26]. The equation is the second member in the entire KP hierarchy [25], originally defined by a Hirota bilinear equation

$$\begin{aligned} B_{JM}(f) &:= (D_x^3 D_y + 2D_t D_y - 3D_x D_z) f \cdot f \\ &= 2(f_{xxx} f - f_y f_{xxx} - 3f_x f_{xy} + 3f_{xx} f_{xy} \\ &\quad + 2f_{yt} f - 2f_y f_t - 3f_{xz} f + 3f_x f_z) = 0, \end{aligned} \quad (4)$$

under the link from f to u :

$$u = 2(\ln f)_x. \quad (5)$$

This link is also a characteristic transformation used in Bell polynomial theories of soliton equations [27, 28] and it actually presents

$$P_{JM}(u) = \frac{f(B_{JM}(f))_x - 2f_x B_{JM}(f)}{f^3}. \quad (6)$$

Therefore, if f solves the bilinear Jimbo-Miwa equation (4), then $u = 2(\ln f)_x$ will engender a solution to the Jimbo-Miwa equation (3).

It is recognized that the Jimbo-Miwa equation (3) passes the Painlevé test only for a subclass of solutions [26] and does not possess a Kac–Moody–Virasoro symmetry algebra [29]. Nevertheless, different types of exact solutions to the Jimbo-Miwa equation (3) are found (see, e.g., [30]–[32]). The Hirota perturbation technique generates one- and two-soliton solutions [26] and dromion-type solutions [32], and the transformed rational function algorithm yields various traveling wave solutions [33]. It is obvious that the Jimbo-Miwa equation (3) has the following y - or x -independent solutions:

$$u = F(x, t) + H(z, t), \quad u = G(y, z) + H(z, t), \quad (7)$$

where F, G and H are arbitrary functions in the indicated variables. These solutions contain more special solutions: $u = F(x, t)$, $u = G(y, z)$ and $u = H(z, t)$, which are independent of two variables of x, y, z, t . Beginning with such solutions, various variable separated solutions are presented by performing Painlevé analysis and abundant nonlinear coherent structures are explored [34]. Moreover, among polynomial functions with individual degrees of the independent variables less than 2, the Jimbo-Miwa equation (3) has the following polynomial solutions:

$$\begin{aligned} u = & a_0 + a_1 x + a_2 y + a_3 z + a_4 t + a_5 xz \\ & + a_6 xt + a_7 yz + \frac{3}{2} a_5 yt + a_8 zt, \end{aligned} \quad (8)$$

where a_i , $0 \leq i \leq 8$, are arbitrary parameters.

In what follows, we concentrate on presenting lump-type solutions to the (3+1)-dimensional Jimbo-Miwa equation (3) by formulating a problem of searching for positive quadratic function solutions to the bilinear Jimbo-Miwa equation (4).

2.2 Lump-type solutions

We apply the computer algebra system Maple to search for quadratic function solutions to the (3+1)-dimensional bilinear Jimbo-Miwa equation (4). A direct Maple symbolic computation with

$$\begin{aligned} f &= g^2 + h^2 + a_{11}, \quad g = a_1 x + a_2 y + a_3 z + a_4 t + a_5, \\ h &= a_6 x + a_7 y + a_8 z + a_9 t + a_{10}, \end{aligned} \quad (9)$$

generates the following four sets of solutions for the parameters a_i , $1 \leq i \leq 11$:

$$\begin{aligned} & \left\{ a_1 = a_1, a_2 = -\frac{a_6 a_7}{a_1}, a_3 = -\frac{2 a_4 a_6 a_7}{3 a_1^2}, a_4 = a_4, a_5 = a_5, \right. \\ & \quad a_6 = a_6, a_7 = a_7, a_8 = \frac{2 a_4 a_7}{3 a_1}, a_9 = \frac{a_4 a_6}{a_1}, \\ & \quad \left. a_{10} = a_{10}, a_{11} = a_{11} \right\}, \\ & \left\{ a_1 = a_1, a_2 = a_7, a_3 = a_8, a_4 = \frac{3 a_1 a_8}{2 a_7}, a_5 = a_5, \right. \\ & \quad a_6 = -a_1, a_7 = a_7, a_8 = a_8, a_9 = -\frac{3 a_1 a_8}{2 a_7}, \\ & \quad \left. a_{10} = a_{10}, a_{11} = a_{11} \right\}, \\ & \left\{ a_1 = 0, a_2 = a_2, a_3 = -\frac{2 a_2 (-3 a_6^3 a_9 + 2 a_4^2 a_{11})}{9 a_6^4}, \right. \\ & \quad \left. a_4 = a_4, a_5 = a_5, a_6 = a_6, \right. \end{aligned}$$

$$\begin{aligned}
a_7 &= -\frac{2a_2a_4a_{11}}{3a_6^3}, \quad a_8 = -\frac{2a_2a_4(a_6^3 + 2a_9a_{11})}{9a_6^4}, \\
a_9 &= a_9, \quad a_{10} = a_{10}, \quad a_{11} = a_{11}, \\
\left\{ \begin{aligned} a_1 &= a_1, \quad a_2 = a_2, \quad a_3 = \frac{2a_1a_2a_4 - a_1a_7a_9 + a_2a_6a_9 + a_4a_6a_7}{3(a_1^2 + a_6^2)}, \\ a_4 &= a_4, \quad a_5 = a_5, \quad a_6 = a_6, \quad a_7 = a_7, \\ a_8 &= \frac{2a_1a_2a_9 + a_1a_4a_7 - a_2a_4a_6 + a_6a_7a_9}{3(a_1^2 + a_6^2)}, \\ a_9 &= a_9, \quad a_{10} = a_{10}, \\ a_{11} &= -\frac{3}{2} \frac{(a_1^5a_2 + a_1^4a_6a_7 + 2a_1^3a_2a_6^2 + 2a_1^2a_6^3a_7 + a_1a_2a_6^4 + a_6^5a_7)}{(a_1a_9 - a_4a_6)(a_1a_7 - a_2a_6)} \end{aligned} \right\}.
\end{aligned}$$

They lead to four classes of quadratic function solutions to the bilinear Jimbo-Miwa equation (4), and the resulting quadratic function solutions, in turn, yield four classes of lump-type solutions to the (3+1)-dimensional Jimbo-Miwa equation (3) through the transformation (5). We list these classes of lump-type solutions as follows.

The first class of lump-type solutions to the Jimbo-Miwa equation (3) reads

$$u_1 = \frac{4}{f_1} \frac{a_4a_1^2t + a_4a_6^2t + a_1^3x + a_1a_6^2x + a_1^2a_5 + a_1a_6a_{10}}{a_1} \quad (10)$$

with

$$\begin{aligned}
f_1 &= \left(a_4t + a_1x - \frac{a_6a_7y}{a_1} - \frac{2a_4a_6a_7z}{3a_1^2} + a_5 \right)^2 \\
&\quad + \left(\frac{a_4a_6t}{a_1} + a_6x + a_7y + \frac{2a_4a_7z}{3a_1} + a_{10} \right)^2 + a_{11}. \quad (11)
\end{aligned}$$

The second class of lump-type solutions to the Jimbo-Miwa equation (3) reads

$$u_2 = \frac{4}{f_2} \frac{a_1(3a_1a_8t + 2a_1a_7x + a_5a_7 - a_7a_{10})}{a_7} \quad (12)$$

with

$$\begin{aligned}
f_2 &= \left(\frac{3a_1a_8t}{2a_7} + a_1x + a_7y + a_8z + a_5 \right)^2 \\
&\quad + \left(-\frac{3a_1a_8t}{2a_7} - a_1x + a_7y + a_8z + a_{10} \right)^2 + a_{11}. \quad (13)
\end{aligned}$$

The third class of lump-type solutions to the Jimbo-Miwa equation (3) reads

$$\begin{aligned}
u_3 &= \frac{4}{9f_3} \frac{-6a_2a_4a_6^3z - 4a_2a_4a_9a_{11}z + 9a_6^4a_{10}9a_6^4a_9t}{a_6^3} \\
&\quad + \frac{9a_6^4a_9t + 9a_6^5x - 6a_2a_4a_6a_{11}y}{a_6^3} \quad (14)
\end{aligned}$$

with

$$\begin{aligned}
f_3 &= \left[a_4t + a_2y - \frac{2a_2(-3a_6^3a_9 + 2a_4^2a_{11})}{9a_6^4}z + a_5 \right]^2 \\
&\quad + \left[a_9t + a_6x - \frac{2a_2a_4a_{11}y}{3a_6^3} \right. \\
&\quad \left. - \frac{2a_2a_4(3a_6^3 + 2a_9a_{11})}{9a_6^4}z + a_{10} \right]^2 + a_{11}. \quad (15)
\end{aligned}$$

The fourth class of lump-type solutions to the Jimbo-Miwa equation (3) reads

$$\begin{aligned}
u_4 &= \frac{4}{3f_4(a_1^2 + a_6^2)} (3a_1^3a_4t + 3a_1^2a_6a_9t + 3a_1a_4a_6^2t \\
&\quad + 3a_6^3a_9t + 3a_1^4x + 6a_1^2a_6^2x + 3a_6^4x + 3a_1^3a_2y \\
&\quad + 3a_1^2a_6a_7y + 3a_1a_2a_6^2y + 3a_6^3a_7y + 2a_1^2a_2a_4z \\
&\quad - 2a_1^2a_7a_9z + 4a_1a_2a_6a_9z + 4a_1a_4a_6a_7z \\
&\quad - 2a_2a_4a_6^2z + 2a_6^2a_7a_9z + 3a_1^3a_5 + 3a_1^2a_6a_{10} \\
&\quad + 3a_1a_5a_6^2 + 3a_6^3a_{10}) \quad (16)
\end{aligned}$$

with

$$\begin{aligned}
f_4 &= \left(a_4t + a_1x + a_2y \right. \\
&\quad \left. + \frac{2a_1a_2a_4 - a_1a_7a_9 + a_2a_6a_9 + a_4a_6a_7}{3(a_1^2 + a_6^2)}z + a_5 \right)^2 \\
&\quad + \left(a_9t + a_6x + a_7y \right. \\
&\quad \left. + \frac{2a_1a_2a_9 + a_1a_4a_7 - a_2a_4a_6 + a_6a_7a_9}{3(a_1^2 + a_6^2)}z + a_{10} \right)^2 \\
&\quad + \frac{(a_1^5a_2 + a_1^4a_6a_7 + 2a_1^3a_2a_6^2 + 2a_1^2a_6^3a_7 + a_1a_2a_6^4 + a_6^5a_7)}{(a_1a_7 - a_2a_6)(a_1a_9 - a_4a_6)}. \quad (17)
\end{aligned}$$

In the above solutions, all parameters a_i involved are arbitrary provided that the solutions are well defined. The third lump-type solution (14) is similar to the lump solution (2) to the KP equation I, but the other three solutions are more general since the two linear waves involved contain all independent variables x, y, z, t .

The analyticity of the first three solutions is guaranteed if $a_{11} > 0$. In the fourth solution in (16), assume that all parameters a_i involved are positive, i.e., $a_i > 0$ for $1 \leq i \leq 4$ and $6 \leq i \leq 9$. Then if $\frac{a_2}{a_7} \neq \frac{a_4}{a_9}$, there are lump-type solutions when taking

$$\frac{a_1}{a_6} \in \left(\frac{a_2}{a_7}, \frac{a_4}{a_9} \right) \text{ or } \frac{a_1}{a_6} \in \left(\frac{a_4}{a_9}, \frac{a_2}{a_7} \right), \quad (18)$$

which leads to $a_{11} > 0$; and if $\frac{a_2}{a_7} = \frac{a_4}{a_9}$, which leads to $a_{11} < 0$, there are singularities in the resulting solution (noting that in the case of $\frac{a_2}{a_7} = \frac{a_4}{a_9}$, a_{11} is well defined iff $\frac{a_1}{a_6} \neq \frac{a_2}{a_7}$). All the above rational function solutions $u_i \rightarrow 0$, $1 \leq i \leq 4$, when the corresponding sum of squares $g^2 + h^2 \rightarrow \infty$. But

they do not approach zero in all directions in $\mathbb{R}^4$ due to the character of (3+1)-dimensions in the resulting solutions, and so we call them lump-type solutions.

For the fourth lump-type solution, let us now fix

$$a_5 = 1, a_{10} = -1,$$

and choose two particular sets of parameters:

$$a_1 = 3, a_2 = 5, a_4 = 4, a_6 = 3, a_7 = 6, a_9 = 3,$$

and

$$a_1 = 4, a_2 = 3, a_4 = 5, a_6 = 3, a_7 = 2, a_9 = 6,$$

which satisfy the two conditions in eq. (18), respectively. The corresponding two particular analytical lump-type solutions of (16) are given by

$$u_{4,1} = \frac{324(21t + 18x + 33y + 26z)}{g_1} \quad (19)$$

with

$$\begin{aligned} g_1 = & 2025t^2 + 3402tx + 6156ty + 4950tz + 1458x^2 \\ & + 5346xy + 4212xz + 4941y^2 + 7686yz + 3050z^2 \\ & + 162t - 162y + 72z + 144504, \end{aligned} \quad (20)$$

and

$$u_{4,2} = \frac{36(950t + 652x + 450y + 462z + 25)}{g_2} \quad (21)$$

with

$$\begin{aligned} g_2 = & 13725t^2 + 17100tx + 12150ty + 13176tz \\ & + 5625x^2 + 8100xy + 8316xz + 2925y^2 \\ & + 5928yz + 3172z^2 - 450t + 450x \\ & + 450y - 84z + 422325, \end{aligned} \quad (22)$$

respectively.

3 Concluding remarks

Based on the Hirota formulation and by a Maple symbolic computation, we presented four classes of lump-type solutions to the (3+1)-dimensional Jimbo-Miwa equation (3), and analyzed the positivity of the fourth quadratic function solution leading to analytical lump-type solutions, of which two particular analytical lump-type solutions were computed under special choices of parameters involved.

We remark that it would be very interesting to determine conditions under which there exist positive polynomial solutions to a kind of generalized bilinear and trilinear differential equations, as did for resonant solutions in terms of exponential functions [35, 36]. This kind of polynomial solutions will generate analytical lump or lump-type solutions, particularly rogue wave solutions, to the corresponding nonlinear equations through $u = 2(\ln f)_x$ or $u = 2(\ln f)_{xx}$. Higher-order rogue wave solutions will be linked to a wide variety of mathematical topics including generalized Wronskian solutions [37] and generalized Darboux transformations [38]. Multicomponent or higher-order generalizations of lump solutions, especially in (3+1)-dimensional cases and fully discrete cases, would be a good topic for future research, exhibiting more diverse soliton phenomena.

Funding: The work was supported in part by the NNSFC under the grants 11371326, 11271008, 11371086 and 1371361, the Fundamental Research Funds for the Central Universities (2013XK03), the Natural Science Foundation of Shandong Province (Grant No. ZR2013AL016), Zhejiang Innovation Project of China (Grant No. T200905), the First-class Discipline of Universities in Shanghai and the Shanghai Univ. Leading Academic Discipline Project (No. A13-0101-12-004), and Shanghai University of Electric Power.

References

- [1] R. Hirota, The direct method in soliton theory, Cambridge University Press, New York, 2004.
- [2] M. J. Ablowitz and P. A. Clarkson, Solitons, nonlinear evolution equations and inverse scattering, Cambridge University Press, Cambridge, 1991.
- [3] P. J. Caudrey, Memories of Hirota's method: Application to the reduced Maxwell-Bloch system in the early 1970s, Phil. Trans. R. Soc. A **369** (2011), 1215–1227.
- [4] A. Tacchella, On rational solutions of multicomponent and matrix KP hierarchies, J. Geom. Phys. **61** (2011), 1319–1328.
- [5] A. -M. Wazwaz, Two B-type Kadomtsev–Petviashvili equations of (2+1) and (3+1) dimensions: Multiple soliton solutions, rational solutions and periodic solutions, Comput. Fluids. **86** (2013), 357–362.
- [6] S. V. Manakov, V. E. Zakharov, L. A. Bortad and V. B. Matveev, Two-dimensional solitons of the Kadomtsev–Petviashvili equation and their interaction, Phys. Lett. A **63** (1977), 205–206.
- [7] D. J. Kaup, The lump solutions and the Bäcklund transformation for the three-dimensional three-wave resonant interaction, J. Math. Phys. **22** (1981), 1176–1181.
- [8] C. R. Gilson and J. J. C. Nimmo, Lump solutions of the BKP equation, Phys. Lett. A **147** (1990), 472–476.

- [9] J. Satsuma and M. J. Ablowitz, Two-dimensional lumps in nonlinear dispersive systems, *J. Math. Phys.* **20** (1979), 1496–1503.
- [10] K. Imai, Dromion and lump solutions of the Ishimori-I equation, *Prog. Theor. Phys.* **98** (1997), 1013–1023.
- [11] P. Müller, C. Garrett and A. Osborne, Rogue waves, *Oceanography* **18** (2005), 66–75.
- [12] D. R. Solli, C. Ropers, P. Koonath and B. Jalali, Optical rogue waves, *Nature* **450** (2007), 1054–1057.
- [13] W. X. Ma and Y. You, Solving the Korteweg-de Vries equation by its bilinear form: Wronskian solutions, *Trans. Amer. Math. Soc.* **357** (2005), 1753–1778.
- [14] W. X. Ma, C. X. Li and J. S. He, A second Wronskian formulation of the KP equation, *Nonlinear Anal. TMA* **70** (2009), 4245–4258.
- [15] F. Gesztesy and W. Schweiger, Rational KP and mKP-solutions in Wronskian form, *Rep. Math. Phys.* **30** (1991), 205–222.
- [16] W. X. Ma and Y. You, Rational solutions of the Toda lattice equation in Casoratian form, *Chaos Solitons Fract.* **22** (2004), 395–406.
- [17] M. Khalfallah, New exact traveling wave solutions of the (3+1) dimensional Kadomtsev–Petviashvili (KP) equation, *Commun. Nonlinear Sci. Numer. Simul.* **14** (2009), 1169–1175.
- [18] D. I. Sinelshchikov, Comment on: new exact traveling wave solutions of the (3+1)-dimensional Kadomtsev–Petviashvili (KP) equation, *Commun. Nonlinear Sci. Numer. Simul.* **15** (2010), 3235–3236.
- [19] W. X. Ma, Comment on the 3+1 dimensional Kadomtsev–Petviashvili equations, *Commun. Nonlinear Sci. Numer. Simul.* **16** (2011), 2663–2666.
- [20] W. A. Malfliet, Solitary wave solutions of nonlinear wave equations, *Am. J. Phys.* **60** (1992), 650–654.
- [21] M. L. Wang, X. Z. Li and J. L. Zhang, The (G'/G) -expansion method and travelling wave solutions of nonlinear evolution equations in mathematical physics, *Phys. Lett. A* **372** (2008), 417–423.
- [22] W. X. Ma and A. Abdeljabbar, A bilinear Bäcklund transformation of a (3+1)-dimensional generalized KP equation, *Appl. Math. Lett.* **12** (2012), 1500–1504.
- [23] Y. Zhang and W. X. Ma, Rational solutions to a KdV-like equation, *Appl. Math. Comput.* **256** (2015), 252–256.
- [24] W. X. Ma, Generalized bilinear differential equations, *Stud. Nonlinear Sci.* **2** (2011), 140–144.
- [25] M. Jimbo and T. Miwa, Solitons and infinite-dimensional Lie algebras, *Publ. Res. Inst. Math. Sci.* **19** (1983), 943–1001.
- [26] B. Dorizzi, B. Grammaticos, A. Ramani and P. Winternitz, Are all the equations of the Kadomtsev–Petviashvili hierarchy integrable? *J. Math. Phys.* **27** (1986), 2848–2852.
- [27] C. Gilson, F. Lambert, J. Nimmo and R. Willox, On the combinatorics of the Hirota D-operators, *Proc. Roy. Soc. London Ser. A* **452** (1996), 223–234.
- [28] W. X. Ma, Bilinear equations, Bell polynomials and linear superposition principle, *J. Phys. Conf. Ser.* **411** (2013), 012021, 11pp.
- [29] J. Rubin and P. Winternitz, Point symmetries of conditionally integrable nonlinear evolution equations, *J. Math. Phys.* **31** (1990), 2085–2090.
- [30] B. Tian and Y. T. Gao, Beyond travelling waves: A new algorithm for solving nonlinear evolution equations, *Comput. Phys. Commun.* **95** (1996), 139–142.
- [31] M. Senthilvelan, On the extended applications of homogeneous balance method, *Appl. Math. Comput.* **123** (2001), 381–388.
- [32] A. -M. Wazwaz, New solutions of distinct physical structures to high-dimensional nonlinear evolution equations, *Appl. Math. Comput.* **196** (2008), 363–370.
- [33] W. X. Ma and J. -H. Lee, A transformed rational function method and exact solutions to the 3+1 dimensional Jimbo-Miwa equation, *Chaos Solitons Fract.* **42** (2009), 1356–1363.
- [34] X. Y. Tang and Z. F. Liang, Variable separated solutions for the (3+1)-dimensional Jimbo-Miwa equation, *Phys. Lett. A* **351** (2006), 398–402.
- [35] W. X. Ma, Bilinear equations and resonant solutions characterized by Bell polynomials, *Rep. Math. Phys.* **72** (2013), 41–56.
- [36] W. X. Ma, Trilinear equations, Bell polynomials, and resonant solutions, *Front. Math. China* **8** (2013), 1139–1156.
- [37] W. X. Ma, Wronskians, Generalized Wronskians and solutions to the Korteweg-de Vries equation, *Chaos Solitons Fract.* **19** (2004), 163–170.
- [38] B. L. Guo, L. M. Ling and Q. P. Liu, Nonlinear Schrödinger equation: Generalized Darboux transformation and rogue wave solutions, *Phys. Rev. E* **85** (2012), 026607, 9pp.