

An integrable generalization of the Kaup–Newell soliton hierarchy

Wen-Xiu Ma¹, Chang-Guang Shi², Emmanuel A Appiah¹, Chunxia Li³ and Shoufeng Shen⁴

¹ Department of Mathematics and Statistics, University of South Florida, Tampa, FL 33620-5700, USA

² College of Mathematics and Physics, Shanghai University of Electric Power, Shanghai 200090, People's Republic of China

³ School of Mathematical Sciences, Capital Normal University, Beijing 100048, People's Republic of China

⁴ Department of Applied Mathematics, Zhejiang University of Technology, Hangzhou 310023, People's Republic of China

E-mail: mawx@cas.usf.edu

Received 27 February 2014, revised 25 April 2014

Accepted for publication 21 May 2014

Published 26 June 2014

Abstract

A generalization of the Kaup–Newell spectral problem associated with $\mathfrak{sl}(2, \mathbb{R})$ is introduced and the corresponding generalized Kaup–Newell hierarchy of soliton equations is generated. Bi-Hamiltonian structures of the resulting soliton hierarchy, leading to a common recursion operator, are furnished by using the trace identity, and thus, the Liouville integrability is shown for all systems in the new generalized soliton hierarchy. The involved bi-Hamiltonian property is explored by using the computer algebra system Maple.

Keywords: soliton equations, Hamiltonian structure, symmetries and conservation laws

1. Introduction

Soliton hierarchies are generated from matrix spectral problems or Lax pairs associated with matrix loop algebras (see, e.g., [1–5]). Among celebrated examples, with dependent variables less than three, are the Korteweg–de Vries hierarchy [6], the Ablowitz–Kaup–Newell–Segur hierarchy [7], the Dirac hierarchy [8], the Kaup–Newell hierarchy [9], the Wadati–Konno–Ichikawa hierarchy [10] and the Boiti–Pempinelli–Tu hierarchy [11]. They often possess bi-Hamiltonian structures and are Liouville integrable. The associated Hamiltonian structures can be furnished by the variational identity [12, 13], particularly by the trace identity when the underlying matrix loop algebra is semisimple [14].

Let us give an outline of the procedure for constructing soliton hierarchies by zero curvature equations (see, e.g., [14, 15]). The beginning is to take a matrix loop algebra $\tilde{\mathfrak{g}}$, associated with a matrix Lie algebra $\mathfrak{g}$ with the commutator

$$[A, B] = AB - BA, \quad A, B \in \mathfrak{g}, \quad (1.1)$$

and introduce spatial matrix spectral problem

$$\phi_x = U\phi, \quad U = U(u, \lambda) \in \tilde{\mathfrak{g}}, \quad (1.2)$$

where u is the potential vector and λ is the spectral parameter (see, e.g., [16] for examples of block matrix loop algebras). We then find a solution

$$W = W(u, \lambda) = \sum_{i \geq 0} W_{0,i} \lambda^{-i}, \quad W_{0,i} \in \mathfrak{g}, \quad i \geq 0, \quad (1.3)$$

to the stationary zero curvature equation

$$W_x = [U, W]. \quad (1.4)$$

Further, we try to formulate the temporal matrix spectral problems

$$\phi_{t_m} = V^{[m]} \phi = V^{[m]}(u, \lambda) \phi, \quad m \geq 0, \quad (1.5)$$

by introducing the Lax matrices

$$V^{[m]} = V^{[m]}(u, \lambda) = (\lambda^m W)_+ + \Delta_m, \quad \Delta_m \in \tilde{\mathfrak{g}}, \quad m \geq 0,$$

P_+ denoting the polynomial part of P in λ , such that the

compatibility conditions of equations (1.2) and (1.5), i.e., the zero curvature equations

$$U_{t_m} - V_x^{[m]} + [U, V^{[m]}] = 0, \quad m \geq 0, \quad (1.6)$$

will engender a hierarchy of soliton equations:

$$u_{t_m} = K_m(u), \quad m \geq 0. \quad (1.7)$$

Such a soliton hierarchy usually possesses a recursion operator and Hamiltonian structures

$$u_{t_m} = K_m(u) = \Phi^m K_0 = J \frac{\delta \mathcal{H}_m}{\delta u}, \quad m \geq 0. \quad (1.8)$$

The recursion operator Φ (see [17] for definition) is generated from the stationary zero curvature equation (1.4) and the Hamiltonian functionals $\mathcal{H}_m$'s can often be computed through the trace identity [14, 15]:

$$\begin{aligned} \frac{\delta}{\delta u} \int \text{tr} \left(\frac{\partial U}{\partial \lambda} W \right) dx &= \lambda^{-\gamma} \frac{\partial}{\partial \lambda} \lambda^\gamma \text{tr} \left(\frac{\partial U}{\partial u} W \right), \\ \gamma &= -\frac{\lambda}{2} \frac{d}{d\lambda} \ln \left| \text{tr} (W^2) \right|, \end{aligned} \quad (1.9)$$

or generally, the variational identity [12, 13]:

$$\begin{aligned} \frac{\delta}{\delta u} \int \left\langle \frac{\partial U}{\partial \lambda}, W \right\rangle dx &= \lambda^{-\gamma} \frac{\partial}{\partial \lambda} \lambda^\gamma \left\langle \frac{\partial U}{\partial u}, W \right\rangle, \\ \gamma &= -\frac{\lambda}{2} \frac{d}{d\lambda} \ln |\langle W, W \rangle|, \end{aligned} \quad (1.10)$$

where $\langle \cdot, \cdot \rangle$ is a non-degenerate, symmetric and ad-invariant bilinear form on the underlying matrix loop algebra $\tilde{\mathfrak{g}}$ [12, 13].

We shall make use of the three-dimensional special linear Lie algebra $\mathfrak{sl}(2, \mathbb{R})$, consisting of trace-free 2×2 matrices. This Lie algebra is simple and has the basis

$$e_1 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad e_3 = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \quad (1.11)$$

whose commutator relations are

$$[e_1, e_2] = 2e_2, \quad [e_1, e_3] = -2e_3, \quad [e_2, e_3] = e_1.$$

Its derived algebra is itself, and so, it is three-dimensional, too. The matrix loop algebra we shall adopt in what follows is

$\tilde{\mathfrak{sl}}(2, \mathbb{R})$

$$= \left\{ \sum_{i \geq 0} M_i \lambda^{n-i} \mid M_i \in \mathfrak{sl}(2, \mathbb{R}), \quad i \geq 0, \quad n \in \mathbb{Z} \right\}. \quad (1.12)$$

It is the space of all Laurent series in λ with coefficients in $\mathfrak{sl}(2, \mathbb{R})$ and a finite regular part. Particular examples of this matrix loop algebra $\tilde{\mathfrak{sl}}(2, \mathbb{R})$ contain matrices of the form

$$\lambda^m e_1 + \lambda^n e_2 + \lambda^l e_3$$

with arbitrary integers m, n, l . The loop algebra $\tilde{\mathfrak{sl}}(2, \mathbb{R})$ lays a foundation for our study of soliton equations, from which many well-known soliton hierarchies are generated (see, e.g., [6–10]).

In this paper, we would like to consider a generalization of the Kaup–Newell spectral problem, which is associated

with the matrix loop algebra $\tilde{\mathfrak{sl}}(2, \mathbb{R})$, and compute an associated hierarchy of bi-Hamiltonian soliton equations by zero curvature equations. The corresponding bi-Hamiltonian structures will be furnished by applying the trace identity and a common recursion operator will be explicitly presented. All soliton equations in the resulting soliton hierarchy will be shown to be Liouville integrable. A few concluding remarks will round off the paper.

The resulting hierarchy provides another concrete example of soliton hierarchies associated with the matrix loop algebra $\tilde{\mathfrak{sl}}(2, \mathbb{R})$. The generalized Kaup–Newell spectral problem is formulated by adding a multiple of the product of the two potentials to the e_1 -part of the Kaup–Newell spectral problem, and interestingly, the perturbed spectral problem still generates an integrable hierarchy represented by local vector fields.

2. A generalization of the Kaup–Newell hierarchy

To present a generalized Kaup–Newell soliton hierarchy, let us introduce a new matrix spectral problem:

$$\phi_x = U\phi = U(u, \lambda)\phi, \quad u = \begin{bmatrix} p \\ q \end{bmatrix}, \quad \phi = \begin{bmatrix} \phi_1 \\ \phi_2 \end{bmatrix}, \quad (2.1)$$

with the spectral matrix U being chosen as

$$\begin{aligned} U &= (\lambda^2 + \alpha pq)e_1 + \lambda p e_2 + \lambda q e_3 \\ &= \begin{bmatrix} \lambda^2 + \alpha pq & \lambda p \\ \lambda q & -\lambda^2 - \alpha pq \end{bmatrix} \in \tilde{\mathfrak{sl}}(2, \mathbb{R}), \end{aligned} \quad (2.2)$$

where α is an arbitrarily given constant. The case of $\alpha = 0$, where the perturbation αpq becomes zero, reduces to the Kaup–Newell spectral problem [9].

First, we solve the stationary zero curvature equation (1.4). The equation (1.4) reads as follows:

$$\begin{cases} a_x = \lambda(pc - qb), \\ b_x = 2\lambda^2 b - 2\lambda pa + 2apqb, \\ c_x = -2\lambda^2 c + 2\lambda qa - 2apqc, \end{cases} \quad (2.3)$$

if W is assumed to be

$$W = ae_1 + be_2 + ce_3 = \begin{bmatrix} a & b \\ c & -a \end{bmatrix} \in \tilde{\mathfrak{sl}}(2, \mathbb{R}). \quad (2.4)$$

Further let a, b and c possess the Laurent expansions:

$$\begin{aligned} a &= \sum_{i \geq 0} a_i \lambda^{-2i}, \\ b &= \sum_{i \geq 0} b_i \lambda^{-2i-1}, \\ c &= \sum_{i \geq 0} c_i \lambda^{-2i-1}, \end{aligned} \quad (2.5)$$

and take the initial data

$$a_0 = 1, \quad b_0 = p, \quad c_0 = q, \quad (2.6)$$

which are required by the equations on the highest powers of

λ in equation (2.3):

$$a_{0,x} = pc_0 - qb_0, \quad b_0 = pa_0, \quad c_0 = qa_0.$$

Then, the system in equation (2.3) becomes

$$\begin{cases} a_{i+1,x} = -\frac{1}{2}(qb_{i,x} + pc_{i,x}) + apq(qb_i - pc_i), \\ b_{i+1} = \frac{1}{2}b_{i,x} + pa_{i+1} - apqb_i, \\ c_{i+1} = -\frac{1}{2}c_{i,x} + qa_{i+1} - apqc_i, \end{cases} \quad i \geq 0, \quad (2.7)$$

the first of which is because from equation (2.3), we have

$$\begin{aligned} qb_x + pc_x &= q(2\lambda^2 b - 2\lambda pa + 2apqb) + p(-2\lambda^2 c + 2\lambda qa - 2apqc) \\ &= 2\lambda^2(qb - pc) + 2apq(qb - pc) \\ &= -2\lambda a_x + 2apq(qb - pc). \end{aligned}$$

The recursion relations in equation (2.7) also tell that

$$\begin{bmatrix} c_{i+1} \\ b_{i+1} \end{bmatrix} = L \begin{bmatrix} c_i \\ b_i \end{bmatrix}, \quad L = \begin{bmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{bmatrix}, \quad i \geq 0, \quad (2.8)$$

where

$$\begin{cases} L_{11} = -\frac{1}{2}\partial - apq - \frac{1}{2}q\partial^{-1}p\partial - \alpha q\partial^{-1}p^2q, \\ L_{12} = -\frac{1}{2}q\partial^{-1}q\partial + \alpha q\partial^{-1}pq^2, \\ L_{21} = -\frac{1}{2}p\partial^{-1}p\partial - \alpha p\partial^{-1}p^2q, \\ L_{22} = \frac{1}{2}\partial - apq - \frac{1}{2}p\partial^{-1}q\partial + \alpha p\partial^{-1}pq^2. \end{cases} \quad (2.9)$$

While using the above recursion relations in equation (2.7), we impose the following conditions on constants of integration:

$$a_i|_{u=0} = b_i|_{u=0} = c_i|_{u=0} = 0, \quad i \geq 1, \quad (2.10)$$

to determine the sequence of $\{a_i, b_i, c_i | i \geq 1\}$ uniquely. This way, the first two sets can be computed as follows:

$$\begin{aligned} a_1 &= -\frac{1}{2}pq, \quad b_1 = \frac{1}{2}p_x - \frac{1}{2}p^2q - \alpha p^2q, \\ c_1 &= -\frac{1}{2}q_x - \frac{1}{2}pq^2 - \alpha pq^2; \\ a_2 &= \frac{1}{4}pq_x - \frac{1}{4}p_xq + \frac{3}{8}p^2q^2 + \alpha p^2q^2, \\ b_2 &= \frac{1}{4}p_{xx} - \frac{3}{4}pp_xq + \frac{3}{8}p^3q^2 - \frac{3}{2}\alpha pp_xq \\ &\quad - \frac{1}{2}\alpha p^2q_x + \alpha^2 p^3q^2 + \frac{3}{2}\alpha p^3q^2, \\ c_2 &= \frac{1}{4}q_{xx} + \frac{3}{4}pq q_x + \frac{3}{8}p^2q^3 + \frac{1}{2}\alpha p_xq^2 \\ &\quad + \frac{3}{2}\alpha pq q_x + \alpha^2 p^2q^3 + \frac{3}{2}\alpha p^2q^3. \end{aligned}$$

We saw above the localness of the first two sets of $\{a_i, b_i, c_i | i \geq 1\}$. This is not an accident. Actually, we have the following result on the localness of the whole sequence of $\{a_i, b_i, c_i | i \geq 1\}$.

Proposition 2.1: Let a_0, b_0, c_0 be given by equation (2.6). Then all functions $a_i, b_i, c_i, i \geq 1$, determined by equation (2.7) under the conditions in equation (2.10), are differential polynomials in u with respect to x , and so, they are all local.

Proof. First from the stationary zero curvature equation $W_x = [U, W]$, we can compute that

$$\frac{d}{dx} \text{tr}(W^2) = 2 \text{tr}(WW_x) = 2 \text{tr}(W[U, W]) = 0,$$

and so, noting $\text{tr}(W^2) = 2(a^2 + bc)$, we have

$$a^2 + bc = (a^2 + bc)|_{u=0} = 1, \quad (2.11)$$

the last step of which follows from the initial data in equation (2.6). Now through the Laurent expansions in equation (2.5), this gives

$$a_i = -\frac{1}{2} \sum_{k+l=i, k, l \geq 1} a_k a_l - \frac{1}{2} \sum_{k+l=i-1, k, l \geq 0} b_k b_l, \quad i \geq 1. \quad (2.12)$$

Finally, based on this recursion relation in equation (2.12) and the last two recursion relations in equation (2.7), an application of the mathematical induction tells that all functions $a_i, b_i, c_i, i \geq 1$, are differential polynomials in u with respect to x , and thus, they are all local. The proof is finished. $\square$

Now, envisaged by the recursion relations in equation (2.7), we, as usual, introduce

$$\begin{aligned} W^{[m]} &= \lambda(\lambda^{2m+1}W)_+ = (\lambda^{2m+2}W)_+ \\ &\quad - a_{m+1}e_1 \in \widetilde{\mathfrak{sl}}(2, \mathbb{R}), \quad m \geq 0, \end{aligned}$$

and consequently, we have

$$\begin{aligned} W_x^{[m]} &= [U, W^{[m]}] \\ &= \begin{bmatrix} 0 & \lambda(b_{m,x} - 2apqb_m) \\ \lambda(c_{m,x} + 2apqc_m) & 0 \end{bmatrix}, \quad m \geq 0. \end{aligned}$$

This is not the same type matrix as the Gateaux derivative operator U' , and so, noting

$$[U, e_1] = -2\lambda p e_2 + 2\lambda q e_3,$$

we take a sequence of Lax operators with modification terms:

$$\begin{aligned} V^{[m]} &= (\lambda^{2m+2}W)_+ + \Delta_m \\ &= W^{[m]} + \delta_m e_1 \in \widetilde{\mathfrak{sl}}(2, \mathbb{R}), \quad m \geq 0, \end{aligned} \quad (2.13)$$

where $\delta_m, m \geq 0$, are functions to be determined and

$\Delta_m = (\delta_m - a_{m+1})e_1$, $m \geq 0$. Then, we find that

$$V_x^{[m]} - [U, V^{[m]}] = \begin{bmatrix} \delta_{m,x} & \lambda(b_{m,x} - 2\alpha p q b_m + 2p\delta_m) \\ \lambda(c_{m,x} + 2\alpha p q c_m - 2q\delta_m) & -\delta_{m,x} \end{bmatrix},$$

$$m \geq 0,$$

and therefore, the corresponding zero curvature

$$U_m - V_x^{[m]} + [U, V^{[m]}] = 0, \quad m \geq 0, \quad (2.14)$$

equivalently engender a hierarchy of equations:

$$\begin{cases} p_m = b_{m,x} - 2\alpha p q b_m + 2p\delta_m, \\ q_m = c_{m,x} + 2\alpha p q c_m - 2q\delta_m, \quad m \geq 0. \\ \alpha(pq)_m = \delta_{m,x}, \end{cases} \quad (2.15)$$

To make the third equation to be compatible with the other two equations in the above system in equation (2.15), we observe that

$$\begin{aligned} \delta_{m,x} &= \alpha(p_m q + p q_m) \\ &= \alpha[q(b_{m,x} - 2\alpha p q b_m + 2p\delta_m) \\ &\quad + p(c_{m,x} + 2\alpha p q c_m - 2q\delta_m)] \\ &= \alpha[q(b_{m,x} - 2\alpha p q b_m) + p(c_{m,x} + 2\alpha p q c_m)] \\ &= -2\alpha a_{m+1,x}, \quad m \geq 0, \end{aligned}$$

the last equality of which is a consequence of equation (2.7). Hence, we can take

$$\delta_m = -2\alpha a_{m+1}, \quad m \geq 0, \quad (2.16)$$

and then from the zero curvature equations (2.14), we obtain a hierarchy of soliton equations

$$u_{t_m} = K_m = \begin{bmatrix} b_{m,x} - 2\alpha p q b_m - 4\alpha p a_{m+1} \\ c_{m,x} + 2\alpha p q c_m + 4\alpha q a_{m+1} \end{bmatrix}, \quad m \geq 0. \quad (2.17)$$

All equations in this hierarchy are local since all functions a_i, b_i, c_i , $i \geq 1$, are differential polynomials in u . The first nonlinear system in the soliton hierarchy equation (2.17) is as follows:

$$u_{t_1} = \begin{bmatrix} p \\ q \end{bmatrix}_{t_1} = K_1$$

$$= \begin{bmatrix} -\frac{1}{2}p_{xx} - pp_x q - \frac{1}{2}p^2 q_x - 2\alpha p p_x q \\ -2\alpha p^2 q_x - 2\alpha^2 p^3 q^2 - \frac{1}{2}\alpha p^3 q^2 \\ -\frac{1}{2}q_{xx} - p q q_x - \frac{1}{2}p_x q^2 - 2\alpha p_x q^2 \\ -2\alpha p q q_x + 2\alpha^2 p^2 q^3 + \frac{1}{2}\alpha p^2 q^3 \end{bmatrix}. \quad (2.18)$$

3. Liouville integrability

We shall show that all systems in the generalized Kaup–Newell soliton hierarchy in equation (2.17) are Liouville integrable (see [18] for definition).

To this end, let us first furnish Hamiltonian structures for the hierarchy in equation (2.17). We shall use the trace identity in equation (1.9) (or generally, the variational identity in equation (1.10)). It is direct to see that

$$\begin{aligned} \frac{\partial U}{\partial \lambda} &= \begin{bmatrix} 2\lambda & p \\ q & -2\lambda \end{bmatrix}, \\ \frac{\partial U}{\partial p} &= \begin{bmatrix} \alpha q & \lambda \\ 0 & -\alpha q \end{bmatrix}, \\ \frac{\partial U}{\partial q} &= \begin{bmatrix} \alpha p & 0 \\ \lambda & -\alpha p \end{bmatrix}, \end{aligned}$$

and so, we have

$$\begin{aligned} \text{tr}\left(W \frac{\partial U}{\partial \lambda}\right) &= 4\lambda a + qb + pc, \\ \text{tr}\left(W \frac{\partial U}{\partial p}\right) &= 2\alpha qa + \lambda c, \\ \text{tr}\left(W \frac{\partial U}{\partial q}\right) &= 2\alpha pa + \lambda b. \end{aligned}$$

Now, the trace identity equation (1.9) in this situation gives

$$\begin{aligned} \frac{\delta}{\delta u} \int (4\lambda a + qb + pc) dx \\ = \lambda^{-\gamma} \frac{\partial}{\partial \lambda} \lambda^\gamma \begin{bmatrix} 2\alpha qa + \lambda c \\ 2\alpha pa + \lambda b \end{bmatrix}. \end{aligned} \quad (3.1)$$

Balancing coefficients of λ^{-2m-1} , $m \geq 0$, in the equality leads to

$$\begin{aligned} \frac{\delta}{\delta u} \int (4a_{m+1} + qb_m + pc_m) dx \\ = (\gamma - 2m) \begin{bmatrix} 2\alpha qa_m + c_m \\ 2\alpha pa_m + b_m \end{bmatrix}, \quad m \geq 0. \end{aligned}$$

The identity with $m = 1$ tells $\gamma = 0$, and thus, we obtain

$$\frac{\delta}{\delta u} \mathcal{H}_m = \begin{bmatrix} 2\alpha qa_m + c_m \\ 2\alpha pa_m + b_m \end{bmatrix}, \quad m \geq 0, \quad (3.2)$$

with the Hamiltonian functionals being defined by

$$\begin{aligned} \mathcal{H}_0 &= \int (1 + 2\alpha) pq \, dx, \\ \mathcal{H}_m &= \int \left(-\frac{4a_{m+1} + qb_m + pc_m}{2m} \right) dx, \quad m \geq 1, \end{aligned} \quad (3.3)$$

the first functional $\mathcal{H}_0$ of which was determined directly from the vector $(2\alpha qa_0 + c_0, 2\alpha pa_0 + b_0)^T$.

Now, noting that

$$\begin{bmatrix} c_m \\ b_m \end{bmatrix} = R \begin{bmatrix} 2\alpha q a_m + c_m \\ 2\alpha p a_m + b_m \end{bmatrix},$$

$$R = \begin{bmatrix} 1 - 2\alpha q \partial^{-1} p & 2\alpha q \partial^{-1} q \\ -2\alpha p \partial^{-1} p & 1 + 2\alpha p \partial^{-1} q \end{bmatrix}, \quad m \geq 0, \quad (3.4)$$

$$\begin{cases} p_{t_m} = 2b_{m+1} - 2pa_{m+1} - 4\alpha pa_{m+1}, \\ q_{t_m} = -2c_{m+1} + 2qa_{m+1} + 4\alpha qa_{m+1}, \end{cases} \quad m \geq 0, \quad (3.5)$$

and

$$a_{m+1} = \partial^{-1}(pc_{m+1} - qb_{m+1}), \quad m \geq 0, \quad (3.6)$$

we find that

$$K_m = \begin{bmatrix} -2(1+2\alpha)p\partial^{-1}p & 2+2(1+2\alpha)p\partial^{-1}q \\ -2+2(1+2\alpha)q\partial^{-1}p & -2(1+2\alpha)q\partial^{-1}q \end{bmatrix} \begin{bmatrix} c_{m+1} \\ b_{m+1} \end{bmatrix}$$

$$= J \begin{bmatrix} 2\alpha qa_{m+1} + c_{m+1} \\ 2\alpha pa_{m+1} + b_{m+1} \end{bmatrix}, \quad m \geq 0. \quad (3.7)$$

with

$$J = \begin{bmatrix} -2(1+4\alpha)p\partial^{-1}p & 2+2(1+4\alpha)p\partial^{-1}q \\ -2+2(1+4\alpha)q\partial^{-1}p & -2(1+4\alpha)q\partial^{-1}q \end{bmatrix}, \quad (3.8)$$

which is a Hamiltonian operator [19]. It follows then that the soliton hierarchy in equation (2.17) has the Hamiltonian structures:

$$u_{t_m} = K_m = J \frac{\delta \mathcal{H}_{m+1}}{\delta u}, \quad m \geq 0, \quad (3.9)$$

where the Hamiltonian operator is defined by equation (3.8) and the Hamiltonian functionals $\mathcal{H}_m$'s are given by equation (3.3).

The resulting functionals can generate infinitely many conservation laws for each soliton system in the soliton hierarchy in equation (2.17). We remark that differential polynomial type conservation laws can also be computed either by computer algebra codes (see, e.g., [20]) or from some Riccati equation associated with the underlying matrix spectral problem (see, e.g., [21–25]).

Obviously, we have

$$\frac{\delta \mathcal{H}_{m+1}}{\delta u} = \Psi \frac{\delta \mathcal{H}_m}{\delta u}, \quad \Psi = R^{-1}LR, \quad (3.10)$$

where the inverse operator of R (see [19, 26] for other applications of such an inverse operator) can be computed as follows:

$$R^{-1} = \begin{bmatrix} 1 + 2\alpha q \partial^{-1} p & -2\alpha q \partial^{-1} q \\ 2\alpha p \partial^{-1} p & 1 - 2\alpha p \partial^{-1} q \end{bmatrix}. \quad (3.11)$$

Then, from $K_{m+1} = \Phi K_m$, $m \geq 0$, and $J\Psi = \Phi J$, we obtain a common recursion operator for the soliton hierarchy in

equation (2.17):

$$\Phi = \Psi^\dagger = R^\dagger L^\dagger (R^{-1})^\dagger,$$

where $Q^\dagger$ denotes the adjoint operator of Q . This recursion operator can be explicitly computed as follows:

$$\Phi = \begin{bmatrix} \Phi_{11} & \Phi_{12} \\ \Phi_{21} & \Phi_{22} \end{bmatrix}, \quad (3.12)$$

where

$$\left\{ \begin{aligned} \Phi_{11} &= \frac{1}{2}\partial - \alpha p q + \alpha p \partial^{-1} q \partial - \frac{1}{2}(1+2\alpha)\partial p \partial^{-1} q \\ &\quad + \alpha(1+2\alpha)p^2 q \partial^{-1} q \\ &\quad - 2\alpha^2 p \partial^{-1} p q^2 - \alpha(1+2\alpha)p \partial^{-1} q \partial p \partial^{-1} q \\ &\quad - \alpha(1+2\alpha)p \partial^{-1} p \partial q \partial^{-1} q, \\ \Phi_{12} &= -\alpha p \partial^{-1} p \partial - \frac{1}{2}(1+2\alpha)\partial p \partial^{-1} p \\ &\quad + \alpha(1+2\alpha)p^2 q \partial^{-1} p - 2\alpha^2 p \partial^{-1} p^2 q \\ &\quad - \alpha(1+2\alpha)p \partial^{-1} q \partial p \partial^{-1} p \\ &\quad - \alpha(1+2\alpha)p \partial^{-1} p \partial q \partial^{-1} p, \\ \Phi_{21} &= -\alpha q \partial^{-1} q \partial - \frac{1}{2}(1+2\alpha)\partial q \partial^{-1} q \\ &\quad - \alpha(1+2\alpha)p q^2 \partial^{-1} q + 2\alpha^2 q \partial^{-1} p q^2 \\ &\quad + \alpha(1+2\alpha)q \partial^{-1} q \partial p \partial^{-1} q \\ &\quad + \alpha(1+2\alpha)q \partial^{-1} p \partial q \partial^{-1} q, \\ \Phi_{22} &= -\frac{1}{2}\partial - \alpha p q + \alpha q \partial^{-1} p \partial - \frac{1}{2}(1+2\alpha)\partial q \partial^{-1} p \\ &\quad - \alpha(1+2\alpha)p q^2 \partial^{-1} p \\ &\quad + 2\alpha^2 q \partial^{-1} p^2 q + \alpha(1+2\alpha)q \partial^{-1} q \partial p \partial^{-1} p \\ &\quad + \alpha(1+2\alpha)q \partial^{-1} p \partial q \partial^{-1} p. \end{aligned} \right. \quad (3.13)$$

Various recursion operators can be found through Lax representations or by computer algebra codes for nonlinear partial differential (see, e.g., [27, 28]). There are also direct algorithms for computing symmetries of nonlinear systems of differential equations (see, e.g., [29]).

It is now a straightforward computation to show that all members in the soliton hierarchy in equation (2.17) are bi-Hamiltonian:

$$u_{t_m} = K_m = J \frac{\delta \mathcal{H}_{m+1}}{\delta u} = M \frac{\delta \mathcal{H}_m}{\delta u}, \quad m \geq 0, \quad (3.14)$$

where the second Hamiltonian operator (see [30, 31] for a bi-Hamiltonian theory) reads

$$M = \Phi J = \begin{bmatrix} M_{11} & M_{12} \\ M_{21} & M_{22} \end{bmatrix}, \quad (3.15)$$

with

$$\left\{ \begin{aligned} M_{11} &= 2\alpha p \partial^{-1} p \partial - 2\alpha \partial p \partial^{-1} p \\ &\quad + 4\alpha^2 p^2 q \partial^{-1} p + 4\alpha^2 p \partial^{-1} p^2 q \\ &\quad - 4\alpha^2 p \partial^{-1} p \partial q \partial^{-1} p - 4\alpha^2 p \partial^{-1} q \partial p \partial^{-1} p, \\ M_{12} &= \partial - 2\alpha p q + 2\alpha \partial p \partial^{-1} q + 2\alpha p \partial^{-1} q \partial \\ &\quad - 4\alpha^2 p \partial^{-1} p q^2 \\ &\quad - 4\alpha^2 p^2 q \partial^{-1} q + 4\alpha^2 p \partial^{-1} p \partial q \partial^{-1} q \\ &\quad + 4\alpha^2 p \partial^{-1} q \partial p \partial^{-1} q, \\ M_{21} &= \partial + 2\alpha p q - 2\alpha q \partial^{-1} p \partial - 2\alpha \partial q \partial^{-1} p \\ &\quad - 4\alpha^2 p q^2 \partial^{-1} p \\ &\quad - 4\alpha^2 q \partial^{-1} p^2 q + 4\alpha^2 q \partial^{-1} p \partial q \partial^{-1} p \\ &\quad + 4\alpha^2 q \partial^{-1} q \partial p \partial^{-1} p, \\ M_{22} &= 2\alpha \partial q \partial^{-1} q - 2\alpha q \partial^{-1} q \partial + 4\alpha^2 p q^2 \partial^{-1} q \\ &\quad + 4\alpha^2 q \partial^{-1} p q^2 \\ &\quad - 4\alpha^2 q \partial^{-1} p \partial q \partial^{-1} q - 4\alpha^2 q \partial^{-1} q \partial p \partial^{-1} q. \end{aligned} \right. \quad (3.16)$$

Thus, we see that the soliton hierarchy in equation (2.17) is Liouville integrable, upon noticing that the vector fields K_n , $n \geq 1$, possess distinct differential orders and that the common conserved functionals $\{\mathcal{H}_n\}_{n=0}^\infty$ and symmetries $\{K_n\}_{n=0}^\infty$ commute:

$$\{\mathcal{H}_k, \mathcal{H}_l\}_J = \int \left(\frac{\delta \mathcal{H}_k}{\delta u} \right)^T J \frac{\delta \mathcal{H}_l}{\delta u} dx = 0, \quad k, l \geq 0, \quad (3.17)$$

$$\{\mathcal{H}_k, \mathcal{H}_l\}_M = \int \left(\frac{\delta \mathcal{H}_k}{\delta u} \right)^T M \frac{\delta \mathcal{H}_l}{\delta u} dx = 0, \quad k, l \geq 0, \quad (3.18)$$

and

$$[K_k, K_l] = K'_k(u)[K_l] - K'_l(u)[K_k] = 0, \quad k, l \geq 0. \quad (3.19)$$

These commuting relations are also consequences of the Virasoro algebra of Lax operators [32–34], which is much easier to be established than a bi-Hamiltonian formulation since it is long to show that J and M constitute a Hamiltonian pair (see [30] for details on the bi-Hamiltonian theory).

4. Concluding remarks

Based on the matrix loop algebra $\tilde{\mathfrak{sl}}(2, \mathbb{R})$, we introduced a generalization of the Kaup–Newell spectral problem, and generated a hierarchy of soliton differential equations from the associated zero curvature equations. The resulting soliton equations are bi-Hamiltonian and Liouville integrable, and their common Hamiltonian pair and recursion operator were explicitly given. The Kaup–Newell soliton hierarchy is the special case with $\alpha = 0$ of the resulting generalized soliton hierarchy.

A hierarchy of generalized AKNS equations was similarly made and its nonlinearization was carried out previously in [35]. Very recently, the special orthogonal Lie algebra $\mathfrak{so}(3, \mathbb{R})$ has also been used to generate new soliton hierarchies

[36–38]. Among typical discussed spectral matrices are the following three:

$$U(u, \lambda) = \lambda e_1 + p e_2 + q e_3,$$

$$U(u, \lambda) = \lambda^2 e_1 + \lambda p e_2 + \lambda q e_3,$$

$$U(u, \lambda) = \lambda e_1 + \lambda p e_2 + \lambda q e_3,$$

where $u = (p, q)^T$ and e_1 , e_2 and e_3 are the basis matrices of $\mathfrak{so}(3, \mathbb{R})$:

$$e_1 = \begin{bmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix},$$

$$e_2 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix},$$

$$e_3 = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

There are many other higher-order matrix spectral problems which lead to soliton hierarchies (see, e.g., [39–44]).

There has also been a growing interest in generating hierarchies of integrable couplings [45] from matrix spectral problems associated with non-semisimple matrix loop algebras [46]. Integrable couplings show very rich structures, bringing us inspiring thoughts and ideas to classify multi-component integrable systems [47]. Bi-integrable couplings and tri-integrable couplings do exhibit diverse nice structures on recursion operators in block matrix form [13, 47]. It should be significantly important to explore more algebraic and geometric mathematical structures on integrable couplings. There are many interesting questions on integrable couplings, which are open to us. For example, is there any Hamiltonian structure for the bi-integrable coupling

$$u_t = K(u), \quad v_t = K'(u)[v], \quad w_t = K'(u)[w],$$

where K' denotes the Gateaux derivative and $u_t = K$ is assumed to be Hamiltonian? How can one generally solve the perturbation system

$$u_t = K(u), \quad v_t = K'(u)[v],$$

even in the KdV case with $K(u) = 6uu_x + u_{xxx}$, which gives

$$u_t = 6uu_x + u_{xxx}, \quad v_t = 6(uv)_x + v_{xxx}?$$

There are, though, plenty of particular solutions to the perturbation system, and one immediate solution is to take v as a symmetry of $u_t = K(u)$.

Acknowledgements

The work was supported in part by NSF under the grant DMS-1301675, NNSFC under the grants 11371326, 11271008, 61072147, 11271266 and 11371323, Natural Science Foundation of Shanghai (Grant No. 11ZR1414100), Zhejiang Innovation Project of China (Grant No. T200905), and the First-class Discipline of Universities in Shanghai and the Shanghai Univ. Leading Academic Discipline Project

(No. A.13-0101-12-004). The authors are also grateful to X Gu, H Y Liu, S Manukure, M Mcanally, S M Yu, W Y Zhang and Y Zhou for their stimulating discussions.

References

- [1] Ablowitz M J and Clarkson P A 1991 *Solitons, Nonlinear Evolution Equations and Inverse Scattering* (Cambridge: Cambridge University Press)
- [2] Novikov S, Manakov S V, Pitaevskii L P and Zakharov V E 1984 *Theory of Solitons—The Inverse Scattering Method*, (New York: Consultants Bureau/A Division of Plenum Publishing Corporation)
- [3] Chen D Y 2006 *Introduction to Solitons*, (Beijing: Science Press)
- [4] Drinfeld V G and Sokolov V V 1981 Equations of Korteweg-de Vries type and simple Lie algebras *Soviet Math. Dokl.* **23** 457–62
- [5] Fokas A S 1987 Symmetries and integrability *Stud. Appl. Math.* **77** 253–99
- [6] Lax P D 1968 Integrals of nonlinear equations of evolution and solitary waves *Commun. Pure Appl. Math.* **21** 467–90
- [7] Ablowitz M J, Kaup D J, Newell A C and Segur H 1974 The inverse scattering transform-Fourier analysis for nonlinear problems *Stud. Appl. Math.* **53** 249–315
- [8] Grosse H 1986 New solitons connected to the Dirac equation *Phys. Rep.* **134** 297–304
- [9] Kaup D J and Newell A C 1978 An exact solution for a derivative nonlinear Schrödinger equation *J. Math. Phys.* **19** 798–801
- [10] Wadati M, Konno K and Ichikawa Y H 1979 New integrable nonlinear evolution equations *J. Phys. Soc. Japan* **47** 1698–700
- [11] Boiti M, Pempinelli F and Tu G Z 1984 Canonical structure of soliton equations via isospectral eigenvalue problems *Nuovo Cimento B* **79** 231–65
- [12] Ma W X and Chen M 2006 Hamiltonian and quasi-Hamiltonian structures associated with semi-direct sums of Lie algebras *J. Phys. A: Math. Gen.* **39** 10787–801
- [13] Ma W X 2010 Variational identities and Hamiltonian structures *Proc. of the 1st Int. Workshop on Nonlinear and Modern Mathl Phys* ed W X Ma, X B Hu and Q P Liu vol 1212 (Melville, NY: American Institute of Physics) pp 1–27
- [14] Tu G Z 1989 On Liouville integrability of zero-curvature equations and the Yang hierarchy *J. Phys. A: Math. Gen.* **22** 2375–92
- [15] Ma W X 1992 A new hierarchy of Liouville integrable generalized Hamiltonian equations and its reduction *Chin. Ann. Math. A* **13** 115–23
- Ma W X 1992 A new hierarchy of Liouville integrable generalized Hamiltonian equations and its reduction *Chin. J. Contemp. Math.* **13** 79–89
- [16] Ma W X 2013 Integrable Couplings and Matrix Loop Algebras *Proc. of the 2nd Int. Workshop on Nonlinear and Modern Mathl Phys* ed W X Ma and D Kaup vol 1562 (Melville, NY: American Institute of Physics) pp 105–22
- [17] Olver P J 1977 Evolution equations possessing infinitely many symmetries *J. Math. Phys.* **18** 1212–5
- [18] Ma W X 2014 An integrable counterpart of the D-AKNS soliton hierarchy from $so(3, \mathbb{R})$ *Phys. Lett. A* **378** 1717–20
- [19] Ma W X and Zhou R G 1999 On inverse recursion operator and tri-Hamiltonian formulation for a Kaup–Newell system of DNLS equations *J. Phys. A: Math. Gen.* **32** L239–42
- [20] Hereman W, Adams P J, Eklund H L, Hickman M S and Herbst B M 2009 Direct methods and symbolic software for conservation laws of nonlinear equations *Advances in Nonlinear Waves and Symbolic Computation, loose errata*, (New York: Nova Science Publishers) pp 19–78
- [21] Miura R M, Gardner C S and Kruskal M D 1968 Korteweg-de Vries equation and generalizations II—Existence of conservation laws and constants of motion *J. Math. Phys.* **9** 1204–9
- [22] Wadati M, Sanuki H and Konno K 1975 Relationships among inverse method Bäcklund transformation and an infinite number of conservation laws *Prog. Theor. Phys.* **53** 419–36
- [23] Alberty T M, Koikawa T and Sasaki R 1982 Canonical structure of soliton equations I *Physica D* **5** 43–65
- [24] Falqui G, Magri F and Pedroni M 1998 Bi-Hamiltonian geometry Darboux coverings, and linearization of the KP hierarchy *Commun. Math. Phys.* **197** 303–24
- [25] Casati P, Della Vedova A and Ortenzi G 2008 The soliton equations associated with the affine Kac–Moody Lie algebra $G_2^{(1)}$ *J. Geom. Phys.* **58** 377–86
- [26] Ma W X and Zhou R G 1999 A coupled AKNS–Kaup–Newell soliton hierarchy *J. Math. Phys.* **40** 4419–28
- [27] Gurses M, Karasu A and Sokolov V 1999 On construction of recursion operators from Lax representation *J. Math. Phys.* **40** 6473–90
- [28] Baldwin D E and Hereman W 2010 A symbolic algorithm for computing recursion operators of nonlinear partial differential equations *Int. J. Comput. Math.* **87** 1094–119
- [29] Fuchssteiner B, Oevel W and Gerdts V (ed) 1999 Algorithms and software for symbolic analysis of nonlinear systems *Mathl. Comput. Modelling* **25** 1–212
- [30] Magri F 1978 A simple model of the integrable Hamiltonian equation *J. Math. Phys.* **19** 1156–62
- [31] Fuchssteiner B and Fokas A S 1981 Symplectic structures, their Bäcklund transformations and hereditary symmetries *Physica D* **4** 47–66
- [32] Ma W X 1992 The algebraic structures of isospectral Lax operators and applications to integrable equations *J. Phys. A: Math. Gen.* **25** 5329–43
- [33] Ma W X 1992 Lax representations and Lax operator algebras of isospectral and nonisospectral hierarchies of evolution equations *J. Math. Phys.* **33** 2464–76
- [34] Ma W X 2013 Lie algebra structures associated with zero curvature equations and generalized zero curvature equations *Br. J. Appl. Sci. Tech.* **3** 1336–44
- [35] Yan Z Y and Zhang H Q 2001 A hierarchy of generalized AKNS equations, N-Hamiltonian structures and finite-dimensional involutive systems and integrable systems *J. Math. Phys.* **42** 330–9
- [36] Ma W X 2013 A soliton hierarchy associated with $so(3, \mathbb{R})$ *Appl. Math. Comput.* **220** 117–22
- [37] Ma W X 2013 A spectral problem based on $so(3, \mathbb{R})$ and its associated commuting soliton equations *J. Math. Phys.* **54** 103509–8
- [38] Ma W X, Manukure S and Zheng H C 2013 A counterpart of the WKI soliton hierarchy associated with $so(3, \mathbb{R})$ *Z. Naturforsch. A* in preparation
- [39] Fordy A P 1984 Derivative nonlinear Schrödinger equations and Hermitian symmetric spaces *J. Phys. A: Math. Gen.* **17** 1235–45
- [40] Ma W X 2001 Binary Bargmann symmetry constraints of soliton equations *Nonlinear Anal. Theory Methods Appl.* **47** 5199–211
- [41] Ma W X and Zhou R G 2002 Adjoint symmetry constraints of multicomponent AKNS equations *Chin. Ann. Math. Ser. B* **23** 373–84
- [42] Li C X 2004 A hierarchy of coupled Korteweg-de Vries equations and the corresponding finite-dimensional integrable system *J. Phys. Soc. Japan* **73** 327–31
- [43] Geng X G and Du D L 2006 Two hierarchies of new nonlinear evolution equations associated with 3×3 matrix spectral problems *Chaos Solitons Fractals* **29** 1165–72

- [44] Zhang Y F and Tam H M 2010 Four Lie algebras associated with R^6 and their applications *J. Math. Phys.* **51** 093514–30
- [45] Ma W X and Fuchssteiner B 1996 Integrable theory of the perturbation equations *Chaos Solitons Fractals* **7** 1227–50
- [46] Ma W X, Xu X X and Zhang Y F 2006 Semi-direct sums of Lie algebras and continuous integrable couplings *Phys. Lett. A* **351** 125–30
- [47] Ma W X, Meng J H and Zhang H Q 2012 Integrable couplings, variational identities and Hamiltonian formulations *Glob. J. Math. Sci.* **1** 1–17