



Integrable counterparts of the D-Kaup–Newell soliton hierarchy



Chang-Guang Shi^a, Wen-Xiu Ma^{b,*}, Morgan McAnally^b

^a College of Mathematics and Physics, Shanghai University of Electric Power, Shanghai 200090, PR China

^b Department of Mathematics and Statistics, University of South Florida, Tampa, FL 33620-5700, USA

ARTICLE INFO

Keywords:

Zero curvature equation
Soliton hierarchy
Symmetry
Trace identity
Hamiltonian structure

ABSTRACT

Two integrable counterparts of the D-Kaup–Newell soliton hierarchy are constructed from a matrix spectral problem associated with the three dimensional special orthogonal Lie algebra $so(3, \mathbb{R})$. An application of the trace identity presents Hamiltonian or quasi-Hamiltonian structures of the resulting counterpart soliton hierarchies, thereby showing their Liouville integrability, i.e., the existence of infinitely many commuting symmetries and conserved densities. The involved Hamiltonian and quasi-Hamiltonian properties are shown by computer algebra systems.

© 2014 Elsevier Inc. All rights reserved.

1. Introduction

Soliton equations generated from zero curvature equations provide concrete examples of integrable equations possessing Hamiltonian structures (see, e.g., [1–5]). It is a starting point of generating soliton equations to formulate matrix spectral problems (or Lax pairs) associated with given matrix loop algebras. The trace identity [6] and the variational identity [7] offer powerful techniques for furnishing their Hamiltonian structures.

Soliton equations usually come in hierarchies. The zero curvature formulation (see, e.g., [6,8]) provides a practical approach for building soliton hierarchies. Typical examples of soliton hierarchies which fit into this formulation include the Korteweg–de Vries hierarchy [9], the Ablowitz–Kaup–Newell–Segur hierarchy [10], the Dirac hierarchy [11], the Kaup–Newell hierarchy [12] and the Wadati–Konno–Ichikawa hierarchy [13]. Note that those hierarchies only involve one or two dependent variables. The case of three or more dependent variables is highly complicated and needs a considerable investment of time. Integrable couplings are such examples with a large number of dependent variables, possessing triangular forms.

Very recently, the three-dimensional special orthogonal Lie algebra $so(3, \mathbb{R})$ has been used in constructing soliton hierarchies (see, e.g., [14,15]). This simple Lie algebra can be realized through 3×3 skew-symmetric matrices, and thus, it has the following basis

$$e_1 = \begin{bmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix}, \quad e_3 = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad (1.1)$$

whose commutation relations read

$$[e_1, e_2] = e_3, \quad [e_2, e_3] = e_1, \quad [e_3, e_1] = e_2. \quad (1.2)$$

* Corresponding author.

Its derived algebra is the algebra itself, and thus, it is also 3-dimensional. The only other three-dimensional real Lie algebra with a three-dimensional derived algebra is the special linear algebra $\mathfrak{sl}(2, \mathbb{R})$, which has been widely used in studying soliton equations in soliton theory (see, e.g., [9–13]).

The matrix loop algebra we shall adopt in this paper is

$$\widetilde{\mathfrak{so}}(3, \mathbb{R}) = \left\{ \sum_{i \geq 0} M_i \lambda^{n-i} \mid M_i \in \mathfrak{so}(3, \mathbb{R}), i \geq 0, n \in \mathbb{Z} \right\}. \quad (1.3)$$

It is the space of all Laurent series in λ with coefficients in $\mathfrak{so}(3, \mathbb{R})$ and a finite regular part. Particular examples of this loop algebra $\widetilde{\mathfrak{so}}(3, \mathbb{R})$ are linear combinations of the form:

$$p_1(\lambda)e_1 + p_2(\lambda)e_2 + p_3(\lambda)e_3$$

with arbitrary Laurent polynomials p_1, p_2, p_3 in λ , which constitute a Lie subalgebra. Due to the circular commutation relations (1.2), the loop algebra $\widetilde{\mathfrak{so}}(3, \mathbb{R})$ provides a good structural basis for our study of soliton equations possessing Hamiltonian structures and quasi-Hamiltonian structures, and several new soliton hierarchies have been already worked out from $\widetilde{\mathfrak{so}}(3, \mathbb{R})$, indeed (see, e.g., [14–19]).

In this paper, we would like to use $\widetilde{\mathfrak{so}}(3, \mathbb{R})$ to introduce a counterpart matrix spectral problem for the D-Kaup–Newell spectral problem (see [20]), and compute two integrable counterpart hierarchies of the D-Kaup–Newell soliton hierarchy [20] by zero curvature equations. An application of the trace identity will engender Hamiltonian or quasi-Hamiltonian structures for every member in the resulting two counterpart soliton hierarchies, and thus, the counterpart soliton hierarchies are Liouville integrable. The new counterpart hierarchies provide two other interesting examples of soliton hierarchies associated with the matrix loop algebra $\widetilde{\mathfrak{so}}(3, \mathbb{R})$. A few concluding remarks and comments will finish the paper.

2. Integrable counterparts of the D-Kaup–Newell hierarchy

To generate integrable counterparts, associated with $\mathfrak{so}(3, \mathbb{R})$, of the D-Kaup–Newell soliton hierarchy [20], let us introduce a new 3×3 matrix spectral problem:

$$\phi_x = U\phi = U(u, \lambda)\phi, \quad u = \begin{bmatrix} p \\ q \\ r \end{bmatrix}, \quad \phi = \begin{bmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \end{bmatrix}, \quad (2.1)$$

where the spectral matrix U is chosen as

$$U = U(u, \lambda) = (\lambda^2 + r)e_1 + \lambda pe_2 + \lambda qe_3 = \begin{bmatrix} 0 & -\lambda q & -\lambda^2 - r \\ \lambda q & 0 & -\lambda p \\ \lambda^2 + r & \lambda p & 0 \end{bmatrix} \in \widetilde{\mathfrak{so}}(3, \mathbb{R}). \quad (2.2)$$

This linear combination is the selfsame as the D-Kaup–Newell one associated with $\mathfrak{sl}(2, \mathbb{R})$ [20].

Once a matrix spectral problem is chosen, it becomes a standard routine to compute soliton hierarchies from the matrix spectral problem by the zero curvature formulation (see [6,8]). First, we solve the stationary zero curvature equation

$$W_x = [U, W], \quad W \in \widetilde{\mathfrak{so}}(3, \mathbb{R}). \quad (2.3)$$

If we assume W to be

$$W = ae_1 + be_2 + ce_3 = \begin{bmatrix} 0 & -c & -a \\ c & 0 & -b \\ a & b & 0 \end{bmatrix}, \quad (2.4)$$

then the Eq. (2.3) becomes

$$\begin{cases} a_x = \lambda pc - \lambda qb, \\ b_x = -\lambda^2 c - rc + \lambda qa, \\ c_x = -\lambda pa + \lambda^2 b + rb. \end{cases} \quad (2.5)$$

Further, let a, b and c possess the following Laurent expansions in λ :

$$a = \sum_{i \geq 0} a_i \lambda^{-2i}, \quad b = \sum_{i \geq 0} b_i \lambda^{-2i-1}, \quad c = \sum_{i \geq 0} c_i \lambda^{-2i-1}, \quad (2.6)$$

and take the initial data

$$a_0 = 1, \quad b_0 = p, \quad c_0 = q, \quad (2.7)$$

which are required by the equations on the highest powers of λ in (2.5):

$$a_{0,x} = pc_0 - qb_0, \quad b_0 = pa_0, \quad c_0 = qa_0.$$

Then, the system (2.5) leads equivalently to

$$\begin{cases} a_{i+1,x} = -pb_{i,x} - qc_{i,x} - r(pc_i - qb_i), \\ b_{i+1} = c_{i,x} + pa_{i+1} - rb_i, \\ c_{i+1} = -b_{i,x} + qa_{i+1} - rc_i, \end{cases} \quad i \geq 0, \quad (2.8)$$

the first of which is because from (2.5), we have

$$-\lambda a_x = -\lambda^2(pc - qb) = p(b_x + rc - \lambda qa) + q(c_x + \lambda pa - rb) = pb_x + qc_x + r(pc - qb).$$

While using the above recursion relations (2.8), we impose the condition that the constants of integration take the value of zero:

$$a_i|_{u=0} = b_i|_{u=0} = c_i|_{u=0} = 0, \quad i \geq 1, \quad (2.9)$$

to determine the sequence of $\{a_i, b_i, c_i | i \geq 1\}$ uniquely. This way, the first two sets can be computed as follows:

$$\begin{aligned} a_1 &= \frac{1}{2}(p^2 + q^2), \quad b_1 = q_x - \frac{1}{2}p(p^2 + q^2) - pr, \quad c_1 = -p_x - \frac{1}{2}(p^2 + q^2)q - qr; \\ a_2 &= p_x q - p q_x + \frac{3}{8}(p^2 + q^2)^2 + (p^2 + q^2)r, \\ b_2 &= -p_{xx} - \frac{3}{2}(p^2 + q^2)q_x - 2q_x r - q r_x + \frac{3}{8}p(p^2 + q^2)^2 + \frac{3}{2}p(p^2 + q^2)r + pr^2, \\ c_2 &= -q_{xx} + \frac{3}{2}p_x(p^2 + q^2) + 2p_x r + pr_x + qr^2 + \frac{3}{2}(p^2 + q^2)qr + \frac{3}{8}(p^2 + q^2)^2 q. \end{aligned}$$

We point out that the localness of the first two sets of $\{a_i, b_i, c_i | i \geq 1\}$ is not an accident, and actually, the functions a_i, b_i, c_i , $i \geq 1$, are all local. We state and prove this localness property as follows.

Proposition 1. The functions a_i, b_i, c_i , $i \geq 1$, recursively defined by (2.8) from the initial values of (2.7), are all local.

Proof. Since the condition (2.9) does not create any nonlocalness problem, we can assume that this condition (2.9) holds, that is, the constants of integration take the value of zero, as we need in our discussion.

First from $W_x = [U, W]$, where W is defined by (2.4) with (2.6), we can have

$$\frac{d}{dx} \text{tr}(W^2) = 2\text{tr}(WW_x) = 2\text{tr}(W[U, W]) = 0,$$

and so, due to $\text{tr}(W^2) = -2(a^2 + b^2 + c^2)$, we arrive at

$$a^2 + b^2 + c^2 = (a^2 + b^2 + c^2)|_{u=0} = 1,$$

the last step of which follows from the initial data (2.7) and the condition (2.9). Then, this implies, by using (2.6), that

$$a_i = -\frac{1}{2} \sum_{k+l=i, k,l \geq 1} a_k a_l - \frac{1}{2} \sum_{k+l=i-1, k,l \geq 0} b_k b_l - \frac{1}{2} \sum_{k+l=i-1, k,l \geq 0} c_k c_l, \quad i \geq 1.$$

Based on this recursion relation and the last two recursion relations in (2.8), an application of the mathematical induction finally shows that all the functions a_i, b_i, c_i , $i \geq 1$, are differential functions in u , and so, they are all local. $\square$

Now as usual, let us calculate

$$(\lambda(\lambda^{2m+1}W)_+)_x - [U, \lambda(\lambda^{2m+1}W)_+] = \begin{bmatrix} 0 & -\lambda(c_{m,x} - rb_m) & 0 \\ \lambda(c_{m,x} - rb_m) & 0 & -\lambda(b_{m,x} + rc_m) \\ 0 & \lambda(b_{m,x} + rc_m) & 0 \end{bmatrix}, \quad m \geq 0,$$

where for a matrix $A = (a_{ij})_{k \times l}$, we defined $(\lambda^m A)_+ = ((\lambda^m a_{ij})_+)_{k \times l}$, in which f_+ denotes the polynomial part of a particular series f :

$$f_+ = \sum_{i=0}^n f_i \lambda^i, \quad \text{when } f = \sum_{i=-\infty}^n f_i \lambda^i, \quad n \geq 0. \quad (2.10)$$

The above matrix is not the same type as the Gateaux derivative matrix U' , and so, we need to introduce modification terms in Lax operators to get soliton equations. In what follows, we shall present two kinds of such modification terms.

Noting

$$[U, e_1] = \lambda q e_2 - \lambda p e_3,$$

we take a sequence of Lax operators with the first kind of modification terms:

$$V_1^{[m]} = \lambda(\lambda^{2m+1}W)_+ + \Delta_{1,m}, \quad \Delta_{1,m} = \beta a_{m+1} e_1 \in \widetilde{\mathfrak{so}}(3, \mathbb{R}), \quad m \geq 0, \quad (2.11)$$

where β is an arbitrarily given constant. Then, we arrive at

$$(V_1^{[m]})_x - [U, V_1^{[m]}] = \begin{bmatrix} 0 & -\lambda(c_{m,x} - rb_m + \beta pa_{m+1}) & -\beta a_{m+1,x} \\ \lambda(c_{m,x} - rb_m + \beta pa_{m+1}) & 0 & -\lambda(b_{m,x} + rc_m - \beta qa_{m+1}) \\ \beta a_{m+1,x} & \lambda(b_{m,x} + rc_m - \beta qa_{m+1}) & 0 \end{bmatrix}, \quad m \geq 0,$$

and further, the corresponding zero curvature equations

$$U_{t_m} - (V_1^{[m]})_x + [U, V_1^{[m]}] = 0, \quad m \geq 0, \quad (2.12)$$

equivalently yield a hierarchy of soliton equations:

$$u_{t_m} = \begin{bmatrix} p \\ q \\ r \end{bmatrix}_{t_m} = K_{1,m} = \begin{bmatrix} b_{m,x} + rc_m - \beta qa_{m+1} \\ c_{m,x} - rb_m + \beta pa_{m+1} \\ \beta a_{m+1,x} \end{bmatrix}, \quad m \geq 0. \quad (2.13)$$

Every system in this counterpart soliton hierarchy is local, since the functions a_i, b_i, c_i , $i \geq 1$, are all local. The first two systems in the counterpart hierarchy (2.13) read

$$\begin{cases} p_{t_0} = p_x + qr + \frac{1}{2}\beta(p^2 + q^2)q, \\ q_{t_0} = q_x - pr - \frac{1}{2}\beta p(p^2 + q^2), \\ r_{t_0} = -\beta(pp_x + qq_x), \end{cases} \quad (2.14)$$

and

$$\begin{cases} p_{t_1} = q_{xx} - \frac{3}{2}p^2p_x - \frac{1}{2}p_xq^2 - pq_xq - 2p_xr - pr_x - qr^2 - \frac{1}{2}(p^2 + q^2)qr - \frac{3}{8}\beta(p^2 + q^2)^2q - \beta(p_xq - pq_x)q - \beta(p^2 + q^2)qr, \\ q_{t_1} = -p_{xx} - 2q_xr - qr_x - \frac{1}{2}p^2q_x - p_xpq - \frac{3}{2}q_xq^2 + \frac{1}{2}p(p^2 + q^2)r + pr^2 + \beta p(p_xq - pq_x) + \beta p(p^2 + q^2)r + \frac{3}{8}\beta p(p^2 + q^2)^2, \\ r_{t_1} = \beta[p_{xx}q - pq_{xx} + \frac{3}{2}(p_xp + q_xq)(p^2 + q^2) + 2(p_xp + q_xq)r + (p^2 + q^2)r_x]. \end{cases} \quad (2.15)$$

Let us now take a sequence of Lax operators with the second kind of modification terms:

$$V_2^{[m]} = \lambda(\lambda^{2m+1}W)_+ + \Delta_{2,m}, \quad \Delta_{2,m} = \beta a_m e_1 \in \widetilde{\mathfrak{so}}(3, \mathbb{R}), \quad m \geq 0, \quad (2.16)$$

where β is an arbitrarily given constant. Similarly, we obtain

$$(V_2^{[m]})_x - [U, V_2^{[m]}] = \begin{bmatrix} 0 & -\lambda(c_{m,x} - rb_m + \beta pa_m) & -\beta a_{m,x} \\ \lambda(c_{m,x} - rb_m + \beta pa_m) & 0 & -\lambda(b_{m,x} + rc_m - \beta qa_m) \\ \beta a_{m,x} & \lambda(b_{m,x} + rc_m - \beta qa_m) & 0 \end{bmatrix}, \quad m \geq 0,$$

and thus, the corresponding zero curvature equations

$$U_{t_m} - (V_2^{[m]})_x + [U, V_2^{[m]}] = 0, \quad m \geq 0, \quad (2.17)$$

present a second hierarchy of soliton equations:

$$u_{t_m} = \begin{bmatrix} p \\ q \\ r \end{bmatrix}_{t_m} = K_{2,m} = \begin{bmatrix} b_{m,x} + rc_m - \beta qa_m \\ c_{m,x} - rb_m + \beta pa_m \\ \beta a_{m,x} \end{bmatrix}, \quad m \geq 0, \quad (2.18)$$

where every member is local. The first two systems in the counterpart hierarchy (2.18) are

$$\begin{cases} p_{t_0} = p_x + qr - \beta p, \\ q_{t_0} = q_x - pr + \beta p, \\ r_{t_0} = 0, \end{cases} \quad (2.19)$$

and

$$\begin{cases} p_{t_1} = q_{xx} - \frac{1}{2}p_x(p^2 + q^2) - p(pp_x + qq_x) - 2p_xr - pr_x - \frac{1}{2}(p^2 + q^2)qr - qr^2 - \frac{1}{2}\beta q(p^2 + q^2), \\ q_{t_1} = -p_{xx} - 2q_xr - qr_x - (pp_x + qq_x)q - \frac{1}{2}(p^2 + q^2)q_x + \frac{1}{2}p(p^2 + q^2)r + pr^2 + \frac{1}{2}\beta p(p^2 + q^2), \\ r_{t_1} = \beta(p_xp + qq_x). \end{cases} \quad (2.20)$$

3. Hamiltonian and quasi-Hamiltonian structures

We shall show that all systems in the counterpart soliton hierarchies (2.13) and (2.18) are Liouville integrable. Let us first establish Hamiltonian structures for the first counterpart hierarchy (2.13) and quasi-Hamiltonian structures for the second counterpart hierarchy (2.18).

We shall use the trace identity [6,8]:

$$\frac{\delta}{\delta u} \int \text{tr} \left(\frac{\partial U}{\partial \lambda} W \right) dx = \lambda^{-\gamma} \frac{\partial}{\partial \lambda} \lambda^\gamma \text{tr} \left(\frac{\partial U}{\partial u} W \right), \quad \gamma = -\frac{\lambda}{2} \frac{d}{d\lambda} \ln |\text{tr}(W^2)|, \quad (3.1)$$

or generally, the variational identity (see [7,21]). It is direct to see that

$$\frac{\partial U}{\partial \lambda} = \begin{bmatrix} 0 & -q & -2\lambda \\ q & 0 & -p \\ 2\lambda & p & 0 \end{bmatrix}, \quad \frac{\partial U}{\partial p} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -\lambda \\ 0 & \lambda & 0 \end{bmatrix}, \quad \frac{\partial U}{\partial q} = \begin{bmatrix} 0 & -\lambda & 0 \\ \lambda & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad \frac{\partial U}{\partial r} = \begin{bmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix},$$

and so, we have

$$\text{tr} \left(W \frac{\partial U}{\partial \lambda} \right) = -4\lambda a - 2qc - 2pb, \quad \text{tr} \left(W \frac{\partial U}{\partial p} \right) = -2\lambda b, \quad \text{tr} \left(W \frac{\partial U}{\partial q} \right) = -2\lambda c, \quad \text{tr} \left(W \frac{\partial U}{\partial r} \right) = -2a.$$

Then, the trace identity (3.1) gives rise to

$$\frac{\delta}{\delta u} \int (2\lambda a + qc + pb) dx = \lambda^{-\gamma} \frac{\partial}{\partial \lambda} \lambda^\gamma \begin{bmatrix} \lambda b \\ \lambda c \\ a \end{bmatrix}.$$

A balance of coefficients of λ^{-2m-1} for each $m \geq 0$ in the equality leads to

$$\frac{\delta}{\delta u} \int (2a_{m+1} + qc_m + pb_m) dx = (\gamma - 2m) \begin{bmatrix} b_m \\ c_m \\ a_m \end{bmatrix}, \quad m \geq 0.$$

The identity with $m = 1$ tells $\gamma = 0$, and thus, we can obtain

$$\frac{\delta}{\delta u} \mathcal{H}_m = \begin{bmatrix} b_m \\ c_m \\ a_m \end{bmatrix}, \quad m \geq 0, \quad (3.2)$$

with the Hamiltonian functionals being defined by

$$\mathcal{H}_0 = \int \left[\frac{1}{2}(p^2 + q^2) + r \right] dx, \quad \mathcal{H}_m = \int \left(-\frac{2a_{m+1} + qc_m + pb_m}{2m} \right) dx, \quad m \geq 1. \quad (3.3)$$

Obviously, it follows from (2.8) that

$$K_{1,m} = \begin{bmatrix} -c_{m+1} + qa_{m+1} - \beta qa_{m+1} \\ b_{m+1} - pa_{m+1} + \beta pa_{m+1} \\ \beta a_{m+1,x} \end{bmatrix} = \begin{bmatrix} -c_{m+1} + (1-\beta)qa_{m+1} \\ b_{m+1} - (1-\beta)pa_{m+1} \\ \beta a_{m+1,x} \end{bmatrix}, \quad m \geq 0,$$

and so, we obtain

$$K_{1,m} = J_1 \begin{bmatrix} b_{m+1} \\ c_{m+1} \\ a_{m+1} \end{bmatrix}, \quad J_1 = \begin{bmatrix} 0 & -1 & (1-\beta)q \\ 1 & 0 & -(1-\beta)p \\ -(1-\beta)q & (1-\beta)p & (2\beta-1)\partial \end{bmatrix}, \quad m \geq 0. \quad (3.4)$$

A direct and easy computation by Maple shows that J_1 is a Hamiltonian operator. It follows now that the first counterpart soliton hierarchy (2.13) has the Hamiltonian structures:

$$u_{t_m} = K_{1,m} = J_1 \frac{\delta \mathcal{H}_{m+1}}{\delta u}, \quad m \geq 0, \quad (3.5)$$

where the Hamiltonian functionals, $\{\mathcal{H}_m\}_{m=0}^\infty$, are given by (3.3) and the Hamiltonian operator J_1 is defined as in (3.4).

Similarly, we can have

$$K_{2,m} = J_2 \begin{bmatrix} b_m \\ c_m \\ a_m \end{bmatrix}, \quad J_2 = \begin{bmatrix} \partial & r & -\beta q \\ -r & \partial & \beta p \\ \beta q & -\beta p & 2\beta \partial \end{bmatrix}, \quad m \geq 0. \quad (3.6)$$

This operator is skew-symmetric but does not satisfy the Jacobi identity, which can be shown by using computer algebra systems. From this, we see that the second counterpart soliton hierarchy (2.18) has the quasi-Hamiltonian structures:

$$u_{t_m} = K_{2,m} = J_2 \frac{\delta \mathcal{H}_m}{\delta u}, \quad m \geq 0, \quad (3.7)$$

where $\mathcal{H}_m$'s are given by (3.3) and J_2 is defined as in (3.6).

The resulting functionals correspond to common conservation laws for each soliton system in the counterpart soliton hierarchy (2.13) or (2.18). We point out that such differential polynomial conservation laws can also be generated either directly by computer algebra codes (see, e.g., [22]) or from some Riccati equation obtained from the underlying matrix spectral problems (see, e.g., [23–25]).

Based on the Hamiltonian structures in (3.5) and the quasi-Hamiltonian structures in (3.7), we can now state that the counterpart soliton hierarchies (2.13) and (2.18) are Liouville integrable, i.e., they possess infinitely many commuting conserved functionals and symmetries:

$$\{\mathcal{H}_k, \mathcal{H}_l\}_{J_i} = \int \left(\frac{\delta \mathcal{H}_k}{\delta u} \right)^T J_i \frac{\delta \mathcal{H}_l}{\delta u} dx = 0, \quad i = 1, 2, \quad k, l \geq 0, \quad (3.8)$$

and

$$[K_{i,k}, K_{i,l}] = K'_{i,k}(u)[K_{i,l}] - K'_{i,l}(u)[K_{i,k}] = 0, \quad i = 1, 2, \quad k, l \geq 0. \quad (3.9)$$

These commuting relations are also consequences of the Virasoro algebra of Lax operators. See [26–28] for a detailed and systematical study on algebraic structures of Lax operators and zero curvature equations.

4. Concluding remarks

Starting with the matrix loop algebra $\widehat{\mathfrak{so}}(3, \mathbb{R})$, we introduced a counterpart matrix spectral problem of the D-Kaup–Newell spectral problem by using the same linear combination of basis matrices, and generated two integrable counterparts of the D-Kaup–Newell soliton hierarchy. All members in the resulting counterpart soliton hierarchies are Hamiltonian or quasi-Hamiltonian, and so, Liouville integrable, i.e., possess infinitely many commuting symmetries and conserved densities.

It is noted that the soliton hierarchy (2.13) is different from a soliton hierarchy presented in [29]. Although the two spectral problems of the two hierarchies look very similar and the two first nonlinear systems are almost the same, it is direct to see that the two hierarchies are very different from each other, if one checks the derivative terms in the two second nonlinear systems. Moreover, we did not find a second class of Hamiltonian structures for our soliton hierarchy (2.13) which needs to be compatible with the presented one (3.5). This is pretty different from the bi-Hamiltonian situation, particularly in the Drinfel'd–Sokolov hierarchies corresponding to the affine twisted Lie algebras [30], and also shows a big difference between our soliton hierarchy (2.13) and the soliton hierarchy by Xia et al. in [29], which was proved to possess a bi-Hamiltonian structure. Quasi-Hamiltonian structures can imply integrability as showed in the recent literature (see, e.g., [7]), but of course, they do not behave like bi-Hamiltonian structures, either.

We point out that among typical discussed spectral matrices associated with $\widehat{\mathfrak{so}}(3, \mathbb{R})$ are the following three:

$$U(u, \lambda) = \lambda e_1 + p e_2 + q e_3,$$

$$U(u, \lambda) = \lambda^2 e_1 + \lambda p e_2 + \lambda q e_3,$$

$$U(u, \lambda) = \lambda e_1 + \lambda p e_2 + \lambda q e_3,$$

where $u = (p, q)^T$ includes two dependent variables. Our examples are two new soliton hierarchies with three dependent variables, fitting into the zero curvature formulation. We hope more examples of such soliton hierarchies with three dependent variables or even more dependent variables, such as four or five dependent variables, can be presented. It is recognized that given a starting matrix loop algebra, it normally needs only a considerable amount of time and computational dexterity to compute hierarchies of integrable equations. Higher-order matrix spectral problems can lead to soliton hierarchies but normally need a more considerable investment of time (see, e.g., [31–36]). Integrable couplings (see, e.g., [37–41]) associated with enlarged matrix loop algebras [42] provide more specific examples of soliton hierarchies generated from higher-order matrix spectral problems. They possess triangular forms [37,21] and their conserved densities can be generated by applying the variational identity [7,21].

Acknowledgements

The work was supported in part by NNSFC under the grants 11371326, 11271008 and 61072147, NSF under the grant DMS-1301675, Natural Science Foundation of Shanghai (Grant No. 11ZR1414100), Zhejiang Innovation Project of China (Grant No. T200905), and the First-class Discipline of Universities in Shanghai and the Shanghai Univ. Leading Academic Discipline Project (No. A.13-0101-12-004). The authors are also grateful to E. A. Appiah, C. X. Li, S. Manukure, S. F. Shen, Y. Q. Yao, S. M. Yu and W. Y. Zhang for their stimulating discussions.

References

- [1] M.J. Ablowitz, P.A. Clarkson, *Solitons, Nonlinear Evolution Equations and Inverse Scattering*, Cambridge University Press, Cambridge, 1991.
- [2] F. Calogero, A. Degasperis, *Spectral Transform and Solitons*, Elsevier, North Holland, 2011.
- [3] S. Novikov, S.V. Manakov, L.P. Pitaevskii, V.E. Zakharov, *Theory of solitons – The inverse scattering method*, Consultants Bureau/A Division of Plenum Publishing Corporation, New York, 1984.
- [4] D.Y. Chen, *Introduction to Solitons*, Science Press, Beijing, 2006.
- [5] V.G. Drinfeld, V.V. Sokolov, Equations of Korteweg–de Vries type and simple Lie algebras, *Soviet Math. Dokl.* 23 (1981) 457–462.
- [6] G.Z. Tu, On Liouville integrability of zero-curvature equations and the Yang hierarchy, *J. Phys. A: Math. Gen.* 22 (1989) 2375–2392.
- [7] W.X. Ma, M. Chen, Hamiltonian and quasi-Hamiltonian structures associated with semi-direct sums of Lie algebras, *J. Phys. A: Math. Gen.* 39 (2006) 10787–10801.
- [8] W.X. Ma, A new hierarchy of Liouville integrable generalized Hamiltonian equations and its reduction, *Chin. Ann. Math. A* 13 (1992) 115–123 (Chn.); W.X. Ma, A new hierarchy of Liouville integrable generalized Hamiltonian equations and its reduction, *Chin. J. Contemp. Math.* 13 (1992) 79–89 (Eng.).
- [9] P.D. Lax, Integrals of nonlinear equations of evolution and solitary waves, *Comm. Pure Appl. Math.* 21 (1968) 467–490.
- [10] M.J. Ablowitz, D.J. Kaup, A.C. Newell, H. Segur, The inverse scattering transform-Fourier analysis for nonlinear problems, *Stud. Appl. Math.* 53 (1974) 249–315.
- [11] H. Grosse, New solitons connected to the Dirac equation, *Phys. Rep.* 134 (1986) 297–304.
- [12] D.J. Kaup, A.C. Newell, An exact solution for a derivative nonlinear Schrödinger equation, *J. Math. Phys.* 19 (1978) 798–801.
- [13] M. Wadati, K. Konno, Y.H. Ichikawa, New integrable nonlinear evolution equations, *J. Phys. Soc. Jpn.* 47 (1979) 1698–1700.
- [14] W.X. Ma, A soliton hierarchy associated with $\mathfrak{so}(3, \mathbb{R})$, *Appl. Math. Comput.* 220 (2013) 117–122.
- [15] W.X. Ma, A spectral problem based on $\mathfrak{so}(3, \mathbb{R})$ and its associated commuting soliton equations, *J. Math. Phys.* 54 (2013) 103509, 8 pp.
- [16] W.X. Ma, S. Manukure, H.C. Zheng, A counterpart of the WKI soliton hierarchy associated with $\mathfrak{so}(3, \mathbb{R})$, *Z. Naturforsch. A* 69 (2014) 411–419.
- [17] W.X. Ma, S.F. Shen, S.M. Yu, H.Q. Zhang, W.Y. Zhang, An integrable $\mathfrak{so}(3, \mathbb{R})$ -counterpart of the Heisenberg soliton hierarchy, *Rep Math Phys.*, to appear.
- [18] C.X. Li, S.F. Shen, W.X. Ma, S.M. Yu, Integrable generalizations of the two new soliton hierarchies of AKNS and KN types associated with $\mathfrak{so}(3, \mathbb{R})$, (preprint, 2014).
- [19] W.X. Ma, An integrable counterpart of the D-AKNS soliton hierarchy from $\mathfrak{so}(3, \mathbb{R})$, *Phys. Lett. A* 378 (2014) 1717–1720.
- [20] X.G. Geng, W.X. Ma, A generalized Kaup–Newell spectral problem, soliton equations and finite-dimensional integrable systems, *Nuovo Cimneto A* 108 (1995) 477–486.
- [21] W.X. Ma, Integrabel couplings and matrix loop algebras, in: W.X. Ma, D. Kaup (Eds.), *Proceedings of the Second International Workshop on Nonlinear and Modern Mathematical Physics*, AIP Conference Proceedings, American Institute of Physics, 1562, Melville, NY, 2013, pp. 105–122.
- [22] W. Hereman, P.J. Adams, H.L. Eklund, M.S. Hickman, B.M. Herbst, Direct methods and symbolic software for conservation laws of nonlinear equations, *Advances in Nonlinear Waves and Symbolic Computation*, loose errata, Nova Sci. Publ., New York, 2009, pp. 19–78.
- [23] R.M. Miura, C.S. Gardner, M.D. Kruskal, Korteweg–de Vries equation and generalizations II - Existence of conservation laws and constants of motion, *J. Math. Phys.* 9 (1968) 1204–1209.
- [24] G. Falqui, F. Magri, M. Pedroni, Bi-Hamiltonian geometry, Darboux coverings, and linearization of the KP hierarchy, *Comm. Math. Phys.* 197 (1998) 303–324.
- [25] P. Casati, A. Della Vedova, G. Ortenzi, The soliton equations associated with the affine Kac–Moody Lie algebra $\mathcal{G}_2^{(1)}$, *J. Geom. Phys.* 58 (2008) 377–386.
- [26] W.X. Ma, The algebraic structures of isospectral Lax operators and applications to integrable equations, *J. Phys. A: Math. Gen.* 25 (1992) 5329–5343.
- [27] W.X. Ma, Lax representations and Lax operator algebras of isospectral and nonisospectral hierarchies of evolution equations, *J. Math. Phys.* 33 (1992) 2464–2476.
- [28] W.X. Ma, Lie algebra structures associated with zero curvature equations and generalized zero curvature equations, *British J. Appl. Sci. Tech.* 3 (2013) 1336–1344.
- [29] T.C. Xia, X.H. Chen, D.Y. Chen, Two types of new Lie algebras and corresponding hierarchies of evolution equations, *Chaos Solitons Fractals* 23 (2005) 1033–1041.
- [30] P. Casati, The bi-Hamiltonian structure of the Drinfel’d–Sokolov hierarchies related to the affine twisted Lie algebras, *Differential Geometry and Applications*, Masaryk University Brno, Czech Republic, 1999, pp. 581–592.
- [31] A.P. Fordy, Derivative nonlinear Schrödinger equations and Hermitian symmetric spaces, *J. Phys. A: Math. Gen.* 17 (1984) 1235–1245.
- [32] W.X. Ma, Binary Bargmann symmetry constraints of soliton equations, *Nonlinear Anal.* 47 (2001) 5199–5211.
- [33] W.X. Ma, R.G. Zhou, Adjoint symmetry constraints of multicomponent AKNS equations, *Chin. Ann. Math. Ser. B* 23 (2002) 373–384.
- [34] C.X. Li, A hierarchy of coupled Korteweg–de Vries equations and the corresponding finite-dimensional integrable system, *J. Phys. Soc. Jpn.* 73 (2004) 327–331.
- [35] X.G. Geng, D.L. Du, Two hierarchies of new nonlinear evolution equations associated with 3×3 matrix spectral problems, *Chaos Solitons Fractals* 29 (2006) 1165–1172.
- [36] Y.F. Zhang, H.W. Tam, Four Lie algebras associated with $\mathfrak{R}^6$ and their applications, *J. Math. Phys.* 51 (2010) 093514, 30 pp.
- [37] W.X. Ma, B. Fuchssteiner, Integrable theory of the perturbation equations, *Chaos Solitons Fractals* 7 (1996) 1227–1250.
- [38] Y.F. Zhang, A generalized multi-component Glachette–Johnson (GJ) hierarchy and its integrable coupling system, *Chaos Solitons Fractals* 21 (2004) 305–310.
- [39] T.C. Xia, F.J. Yu, Y. Zhang, The multi-component coupled Burgers hierarchy of soliton equations and its multi-component integrable couplings system with two arbitrary functions, *Physica A* 343 (2004) 238–246.
- [40] Z. Li, H.H. Dong, Two integrable couplings of the Tu hierarchy and their Hamiltonian structures, *Comput. Math. Appl.* 55 (2008) 2643–2652.
- [41] B.L. Feng, J.Q. Liu, A new Lie algebra along with its induced Lie algebra and associated with applications, *Commun. Nonlinear Sci. Numer. Simul.* 16 (2011) 1734–1741.
- [42] W.X. Ma, X.X. Xu, Y.F. Zhang, Semi-direct sums of Lie algebras and continuous integrable couplings, *Phys. Lett. A* 351 (2006) 125–130.